



United States Patent [19]

Stashenko et al.

[11] Patent Number: 5,552,281
[45] Date of Patent: Sep. 3, 1996

[54] HUMAN OSTEOCLAST-SPECIFIC AND -RELATED GENES

[75] Inventors: Philip Stashenko, Norfolk; Yi-Ping Li, Boston; Anne L. Wucherpfennig, Brookline, all of Mass.

[73] Assignee: Forsyth Dental Infirmary for Children, Boston, Mass.

[21] Appl. No.: 392,678

[22] Filed: Feb. 23, 1995

Related U.S. Application Data

[63] Continuation of Ser. No. 45,270, Apr. 6, 1993, abandoned.

[51] Int. Cl.⁶ C07H 21/04; C12N 5/10; C12N 15/70; C12Q 1/68

[52] U.S. Cl. 435/6; 435/69.1; 435/172.3; 435/252.3; 435/320.1; 536/23.1

[58] Field of Search 435/6, 320.1, 252.3, 435/69.1, 172.3; 536/23.1

[56] References Cited

PUBLICATIONS

Blair, Harry C., et al., "Extracellular-matrix degradation at acid pH. Avian osteoclast acid collagenase isolation and characterization", *Biochemical Journal* 290(3):873-884 (15 Mar. 1993).

Tezuka, Ken-Ichi, et al., "Identification of osteopontin in isolated rabbit osteoclasts", *Biochemical and Biophysical Research Communications* 186(2):914-916 (31 Jul. 1992). Tezuka, Ken-Ichi, et al., "Molecular cloning of a possible cysteine proteinase predominantly expressed in osteoclasts", *Journal of Biological Chemistry* 269(2):1106-1108, (14 Jan. 1994).

Horton, Michael A. et al., "Monoclonal Antibodies to Osteoclastomas (Giant Cell Bone Tumors): Definition of Osteoclast-specific Cellular Antigens," *Cancer Research* 45, 5663-5669 (Nov. 1985).

Davies, John et al., "The Osteoclast Functional Antigen, Implicated in the Regulation of Bone Resorption, Is Biochemically Related to the Vitronectin Receptor," *The Journal of Cell Biology* 109, 1817-1826 (Oct. 1989).

Hayman, Alison, R. et al., "Purification and characterization of a tartrate-resistant acid phosphatase from human osteoclastomas," *Biochem. J.* 261, 601-609 (1989).

Sandberg, M. et al., "Localization of the Expression of Types I, III, and IV Collagen, TGF- β 1 and c-fos Genes in Developing Human Calvarial Bones," *Developmental Biology* 130, 324-334 (1988).

Sandberg, M. et al., "Enhanced expression of TGF- β and c-fos mRNAs in the growth plates of developing human long bones," *Development* 102, 461-470 (1988).

Ek-Rylander, Barbro et al., "Cloning, Sequence, and Developmental Expression of a Type 5, Tartrate-resistant, Acid Phosphatase of Rat Bone," *The Journal of Biological Chemistry* 266(36), 24684-24689 (Dec. 25, 1991).

GenBank/EMBL Sequence Search Printout, pp. 1-19 (Jun. 24, 1993).

Primary Examiner—W. Gary Jones

Assistant Examiner—Paul B. Tran

Attorney, Agent, or Firm—Hamilton, Brook, Smith & Reynolds, P.C.

[57]

ABSTRACT

The present invention relates to purified DNA sequences encoding all or a portion of an osteoclast-specific or -related gene products and a method for identifying such sequences. The invention also relates to antibodies directed against an osteoclast-specific or -related gene product. Also claimed are DNA constructs capable of replicating DNA encoding all or a portion of an osteoclast-specific or -related gene product, and DNA constructs capable of directing expression in a host cell of an osteoclast-specific or -related gene product.

5 Claims, 1 Drawing Sheet

1. MACROPHAGE SECRETED TRACHECTOMY SECRETORY SECRETIN-LIKE PROTEIN
61. SECRETIN-LIKE PROTEIN
121. CYTOKINE ANTI-ADHERENT SECRETIN-LIKE PROTEIN
181. CYTOKINE SECRETIN-LIKE PROTEIN
241. CYTOKINE SECRETIN-LIKE PROTEIN
301. CYTOKINE SECRETIN-LIKE PROTEIN
361. CYTOKINE SECRETIN-LIKE PROTEIN
421. CYTOKINE SECRETIN-LIKE PROTEIN
481. CYTOKINE SECRETIN-LIKE PROTEIN
541. CYTOKINE SECRETIN-LIKE PROTEIN
601. CYTOKINE SECRETIN-LIKE PROTEIN
661. CYTOKINE SECRETIN-LIKE PROTEIN
721. CYTOKINE SECRETIN-LIKE PROTEIN
781. CYTOKINE SECRETIN-LIKE PROTEIN
841. CYTOKINE SECRETIN-LIKE PROTEIN
901. CYTOKINE SECRETIN-LIKE PROTEIN
961. CYTOKINE SECRETIN-LIKE PROTEIN
1021. CYTOKINE SECRETIN-LIKE PROTEIN
1081. CYTOKINE SECRETIN-LIKE PROTEIN
1141. CYTOKINE SECRETIN-LIKE PROTEIN
1201. CYTOKINE SECRETIN-LIKE PROTEIN
1261. CYTOKINE SECRETIN-LIKE PROTEIN
1321. CYTOKINE SECRETIN-LIKE PROTEIN
1381. CYTOKINE SECRETIN-LIKE PROTEIN
1441. CYTOKINE SECRETIN-LIKE PROTEIN
1501. CYTOKINE SECRETIN-LIKE PROTEIN
1561. CYTOKINE SECRETIN-LIKE PROTEIN
1621. CYTOKINE SECRETIN-LIKE PROTEIN
1681. CYTOKINE SECRETIN-LIKE PROTEIN
1741. CYTOKINE SECRETIN-LIKE PROTEIN
1801. CYTOKINE SECRETIN-LIKE PROTEIN
1861. CYTOKINE SECRETIN-LIKE PROTEIN
1921. CYTOKINE SECRETIN-LIKE PROTEIN
1981. CYTOKINE SECRETIN-LIKE PROTEIN
2041. CYTOKINE SECRETIN-LIKE PROTEIN
2101. CYTOKINE SECRETIN-LIKE PROTEIN
2161. CYTOKINE SECRETIN-LIKE PROTEIN
2221. CYTOKINE SECRETIN-LIKE PROTEIN
2281. CYTOKINE SECRETIN-LIKE PROTEIN
2341. CYTOKINE SECRETIN-LIKE PROTEIN
2401. CYTOKINE SECRETIN-LIKE PROTEIN
2461. CYTOKINE SECRETIN-LIKE PROTEIN
2521. CYTOKINE SECRETIN-LIKE PROTEIN
2581. CYTOKINE SECRETIN-LIKE PROTEIN
2641. CYTOKINE SECRETIN-LIKE PROTEIN
2701. CYTOKINE SECRETIN-LIKE PROTEIN
2761. CYTOKINE SECRETIN-LIKE PROTEIN
2821. CYTOKINE SECRETIN-LIKE PROTEIN
2881. CYTOKINE SECRETIN-LIKE PROTEIN
2941. CYTOKINE SECRETIN-LIKE PROTEIN
3001. CYTOKINE SECRETIN-LIKE PROTEIN
3061. CYTOKINE SECRETIN-LIKE PROTEIN
3121. CYTOKINE SECRETIN-LIKE PROTEIN
3181. CYTOKINE SECRETIN-LIKE PROTEIN
3241. CYTOKINE SECRETIN-LIKE PROTEIN
3301. CYTOKINE SECRETIN-LIKE PROTEIN
3361. CYTOKINE SECRETIN-LIKE PROTEIN
3421. CYTOKINE SECRETIN-LIKE PROTEIN
3481. CYTOKINE SECRETIN-LIKE PROTEIN
3541. CYTOKINE SECRETIN-LIKE PROTEIN
3601. CYTOKINE SECRETIN-LIKE PROTEIN
3661. CYTOKINE SECRETIN-LIKE PROTEIN
3721. CYTOKINE SECRETIN-LIKE PROTEIN
3781. CYTOKINE SECRETIN-LIKE PROTEIN
3841. CYTOKINE SECRETIN-LIKE PROTEIN
3901. CYTOKINE SECRETIN-LIKE PROTEIN
3961. CYTOKINE SECRETIN-LIKE PROTEIN
4021. CYTOKINE SECRETIN-LIKE PROTEIN
4081. CYTOKINE SECRETIN-LIKE PROTEIN
4141. CYTOKINE SECRETIN-LIKE PROTEIN
4201. CYTOKINE SECRETIN-LIKE PROTEIN
4261. CYTOKINE SECRETIN-LIKE PROTEIN
4321. CYTOKINE SECRETIN-LIKE PROTEIN
4381. CYTOKINE SECRETIN-LIKE PROTEIN
4441. CYTOKINE SECRETIN-LIKE PROTEIN
4501. CYTOKINE SECRETIN-LIKE PROTEIN
4561. CYTOKINE SECRETIN-LIKE PROTEIN
4621. CYTOKINE SECRETIN-LIKE PROTEIN
4681. CYTOKINE SECRETIN-LIKE PROTEIN
4741. CYTOKINE SECRETIN-LIKE PROTEIN
4801. CYTOKINE SECRETIN-LIKE PROTEIN
4861. CYTOKINE SECRETIN-LIKE PROTEIN
4921. CYTOKINE SECRETIN-LIKE PROTEIN
4981. CYTOKINE SECRETIN-LIKE PROTEIN
5041. CYTOKINE SECRETIN-LIKE PROTEIN
5101. CYTOKINE SECRETIN-LIKE PROTEIN
5161. CYTOKINE SECRETIN-LIKE PROTEIN
5221. CYTOKINE SECRETIN-LIKE PROTEIN
5281. CYTOKINE SECRETIN-LIKE PROTEIN
5341. CYTOKINE SECRETIN-LIKE PROTEIN
5401. CYTOKINE SECRETIN-LIKE PROTEIN
5461. CYTOKINE SECRETIN-LIKE PROTEIN
5521. CYTOKINE SECRETIN-LIKE PROTEIN
5581. CYTOKINE SECRETIN-LIKE PROTEIN
5641. CYTOKINE SECRETIN-LIKE PROTEIN
5701. CYTOKINE SECRETIN-LIKE PROTEIN
5761. CYTOKINE SECRETIN-LIKE PROTEIN
5821. CYTOKINE SECRETIN-LIKE PROTEIN
5881. CYTOKINE SECRETIN-LIKE PROTEIN
5941. CYTOKINE SECRETIN-LIKE PROTEIN
6001. CYTOKINE SECRETIN-LIKE PROTEIN
6061. CYTOKINE SECRETIN-LIKE PROTEIN
6121. CYTOKINE SECRETIN-LIKE PROTEIN
6181. CYTOKINE SECRETIN-LIKE PROTEIN
6241. CYTOKINE SECRETIN-LIKE PROTEIN
6301. CYTOKINE SECRETIN-LIKE PROTEIN
6361. CYTOKINE SECRETIN-LIKE PROTEIN
6421. CYTOKINE SECRETIN-LIKE PROTEIN
6481. CYTOKINE SECRETIN-LIKE PROTEIN
6541. CYTOKINE SECRETIN-LIKE PROTEIN
6601. CYTOKINE SECRETIN-LIKE PROTEIN
6661. CYTOKINE SECRETIN-LIKE PROTEIN
6721. CYTOKINE SECRETIN-LIKE PROTEIN
6781. CYTOKINE SECRETIN-LIKE PROTEIN
6841. CYTOKINE SECRETIN-LIKE PROTEIN
6901. CYTOKINE SECRETIN-LIKE PROTEIN
6961. CYTOKINE SECRETIN-LIKE PROTEIN
7021. CYTOKINE SECRETIN-LIKE PROTEIN
7081. CYTOKINE SECRETIN-LIKE PROTEIN
7141. CYTOKINE SECRETIN-LIKE PROTEIN
7201. CYTOKINE SECRETIN-LIKE PROTEIN
7261. CYTOKINE SECRETIN-LIKE PROTEIN
7321. CYTOKINE SECRETIN-LIKE PROTEIN
7381. CYTOKINE SECRETIN-LIKE PROTEIN
7441. CYTOKINE SECRETIN-LIKE PROTEIN
7501. CYTOKINE SECRETIN-LIKE PROTEIN
7561. CYTOKINE SECRETIN-LIKE PROTEIN
7621. CYTOKINE SECRETIN-LIKE PROTEIN
7681. CYTOKINE SECRETIN-LIKE PROTEIN
7741. CYTOKINE SECRETIN-LIKE PROTEIN
7801. CYTOKINE SECRETIN-LIKE PROTEIN
7861. CYTOKINE SECRETIN-LIKE PROTEIN
7921. CYTOKINE SECRETIN-LIKE PROTEIN
7981. CYTOKINE SECRETIN-LIKE PROTEIN
8041. CYTOKINE SECRETIN-LIKE PROTEIN
8101. CYTOKINE SECRETIN-LIKE PROTEIN
8161. CYTOKINE SECRETIN-LIKE PROTEIN
8221. CYTOKINE SECRETIN-LIKE PROTEIN
8281. CYTOKINE SECRETIN-LIKE PROTEIN
8341. CYTOKINE SECRETIN-LIKE PROTEIN
8401. CYTOKINE SECRETIN-LIKE PROTEIN
8461. CYTOKINE SECRETIN-LIKE PROTEIN
8521. CYTOKINE SECRETIN-LIKE PROTEIN
8581. CYTOKINE SECRETIN-LIKE PROTEIN
8641. CYTOKINE SECRETIN-LIKE PROTEIN
8701. CYTOKINE SECRETIN-LIKE PROTEIN
8761. CYTOKINE SECRETIN-LIKE PROTEIN
8821. CYTOKINE SECRETIN-LIKE PROTEIN
8881. CYTOKINE SECRETIN-LIKE PROTEIN
8941. CYTOKINE SECRETIN-LIKE PROTEIN
9001. CYTOKINE SECRETIN-LIKE PROTEIN
9061. CYTOKINE SECRETIN-LIKE PROTEIN
9121. CYTOKINE SECRETIN-LIKE PROTEIN
9181. CYTOKINE SECRETIN-LIKE PROTEIN
9241. CYTOKINE SECRETIN-LIKE PROTEIN
9301. CYTOKINE SECRETIN-LIKE PROTEIN
9361. CYTOKINE SECRETIN-LIKE PROTEIN
9421. CYTOKINE SECRETIN-LIKE PROTEIN
9481. CYTOKINE SECRETIN-LIKE PROTEIN
9541. CYTOKINE SECRETIN-LIKE PROTEIN
9601. CYTOKINE SECRETIN-LIKE PROTEIN
9661. CYTOKINE SECRETIN-LIKE PROTEIN
9721. CYTOKINE SECRETIN-LIKE PROTEIN
9781. CYTOKINE SECRETIN-LIKE PROTEIN
9841. CYTOKINE SECRETIN-LIKE PROTEIN
9901. CYTOKINE SECRETIN-LIKE PROTEIN
9961. CYTOKINE SECRETIN-LIKE PROTEIN
10021. CYTOKINE SECRETIN-LIKE PROTEIN
10081. CYTOKINE SECRETIN-LIKE PROTEIN
10141. CYTOKINE SECRETIN-LIKE PROTEIN
10201. CYTOKINE SECRETIN-LIKE PROTEIN
10261. CYTOKINE SECRETIN-LIKE PROTEIN
10321. CYTOKINE SECRETIN-LIKE PROTEIN
10381. CYTOKINE SECRETIN-LIKE PROTEIN
10441. CYTOKINE SECRETIN-LIKE PROTEIN
10501. CYTOKINE SECRETIN-LIKE PROTEIN
10561. CYTOKINE SECRETIN-LIKE PROTEIN
10621. CYTOKINE SECRETIN-LIKE PROTEIN
10681. CYTOKINE SECRETIN-LIKE PROTEIN
10741. CYTOKINE SECRETIN-LIKE PROTEIN
10801. CYTOKINE SECRETIN-LIKE PROTEIN
10861. CYTOKINE SECRETIN-LIKE PROTEIN
10921. CYTOKINE SECRETIN-LIKE PROTEIN
10981. CYTOKINE SECRETIN-LIKE PROTEIN
11041. CYTOKINE SECRETIN-LIKE PROTEIN
11101. CYTOKINE SECRETIN-LIKE PROTEIN
11161. CYTOKINE SECRETIN-LIKE PROTEIN
11221. CYTOKINE SECRETIN-LIKE PROTEIN
11281. CYTOKINE SECRETIN-LIKE PROTEIN
11341. CYTOKINE SECRETIN-LIKE PROTEIN
11401. CYTOKINE SECRETIN-LIKE PROTEIN
11461. CYTOKINE SECRETIN-LIKE PROTEIN
11521. CYTOKINE SECRETIN-LIKE PROTEIN
11581. CYTOKINE SECRETIN-LIKE PROTEIN
11641. CYTOKINE SECRETIN-LIKE PROTEIN
11701. CYTOKINE SECRETIN-LIKE PROTEIN
11761. CYTOKINE SECRETIN-LIKE PROTEIN
11821. CYTOKINE SECRETIN-LIKE PROTEIN
11881. CYTOKINE SECRETIN-LIKE PROTEIN
11941. CYTOKINE SECRETIN-LIKE PROTEIN
12001. CYTOKINE SECRETIN-LIKE PROTEIN
12061. CYTOKINE SECRETIN-LIKE PROTEIN
12121. CYTOKINE SECRETIN-LIKE PROTEIN
12181. CYTOKINE SECRETIN-LIKE PROTEIN
12241. CYTOKINE SECRETIN-LIKE PROTEIN
12301. CYTOKINE SECRETIN-LIKE PROTEIN
12361. CYTOKINE SECRETIN-LIKE PROTEIN
12421. CYTOKINE SECRETIN-LIKE PROTEIN
12481. CYTOKINE SECRETIN-LIKE PROTEIN
12541. CYTOKINE SECRETIN-LIKE PROTEIN
12601. CYTOKINE SECRETIN-LIKE PROTEIN
12661. CYTOKINE SECRETIN-LIKE PROTEIN
12721. CYTOKINE SECRETIN-LIKE PROTEIN
12781. CYTOKINE SECRETIN-LIKE PROTEIN
12841. CYTOKINE SECRETIN-LIKE PROTEIN
12901. CYTOKINE SECRETIN-LIKE PROTEIN
12961. CYTOKINE SECRETIN-LIKE PROTEIN
13021. CYTOKINE SECRETIN-LIKE PROTEIN
13081. CYTOKINE SECRETIN-LIKE PROTEIN
13141. CYTOKINE SECRETIN-LIKE PROTEIN
13201. CYTOKINE SECRETIN-LIKE PROTEIN
13261. CYTOKINE SECRETIN-LIKE PROTEIN
13321. CYTOKINE SECRETIN-LIKE PROTEIN
13381. CYTOKINE SECRETIN-LIKE PROTEIN
13441. CYTOKINE SECRETIN-LIKE PROTEIN
13501. CYTOKINE SECRETIN-LIKE PROTEIN
13561. CYTOKINE SECRETIN-LIKE PROTEIN
13621. CYTOKINE SECRETIN-LIKE PROTEIN
13681. CYTOKINE SECRETIN-LIKE PROTEIN
13741. CYTOKINE SECRETIN-LIKE PROTEIN
13801. CYTOKINE SECRETIN-LIKE PROTEIN
13861. CYTOKINE SECRETIN-LIKE PROTEIN
13921. CYTOKINE SECRETIN-LIKE PROTEIN
13981. CYTOKINE SECRETIN-LIKE PROTEIN
14041. CYTOKINE SECRETIN-LIKE PROTEIN
14101. CYTOKINE SECRETIN-LIKE PROTEIN
14161. CYTOKINE SECRETIN-LIKE PROTEIN
14221. CYTOKINE SECRETIN-LIKE PROTEIN
14281. CYTOKINE SECRETIN-LIKE PROTEIN
14341. CYTOKINE SECRETIN-LIKE PROTEIN
14401. CYTOKINE SECRETIN-LIKE PROTEIN
14461. CYTOKINE SECRETIN-LIKE PROTEIN
14521. CYTOKINE SECRETIN-LIKE PROTEIN
14581. CYTOKINE SECRETIN-LIKE PROTEIN
14641. CYTOKINE SECRETIN-LIKE PROTEIN
14701. CYTOKINE SECRETIN-LIKE PROTEIN
14761. CYTOKINE SECRETIN-LIKE PROTEIN
14821. CYTOKINE SECRETIN-LIKE PROTEIN
14881. CYTOKINE SECRETIN-LIKE PROTEIN
14941. CYTOKINE SECRETIN-LIKE PROTEIN
15001. CYTOKINE SECRETIN-LIKE PROTEIN
15061. CYTOKINE SECRETIN-LIKE PROTEIN
15121. CYTOKINE SECRETIN-LIKE PROTEIN
15181. CYTOKINE SECRETIN-LIKE PROTEIN
15241. CYTOKINE SECRETIN-LIKE PROTEIN
15301. CYTOKINE SECRETIN-LIKE PROTEIN
15361. CYTOKINE SECRETIN-LIKE PROTEIN
15421. CYTOKINE SECRETIN-LIKE PROTEIN
15481. CYTOKINE SECRETIN-LIKE PROTEIN
15541. CYTOKINE SECRETIN-LIKE PROTEIN
15601. CYTOKINE SECRETIN-LIKE PROTEIN
15661. CYTOKINE SECRETIN-LIKE PROTEIN
15721. CYTOKINE SECRETIN-LIKE PROTEIN
15781. CYTOKINE SECRETIN-LIKE PROTEIN
15841. CYTOKINE SECRETIN-LIKE PROTEIN
15901. CYTOKINE SECRETIN-LIKE PROTEIN
15961. CYTOKINE SECRETIN-LIKE PROTEIN
16021. CYTOKINE SECRETIN-LIKE PROTEIN
16081. CYTOKINE SECRETIN-LIKE PROTEIN
16141. CYTOKINE SECRETIN-LIKE PROTEIN
16201. CYTOKINE SECRETIN-LIKE PROTEIN
16261. CYTOKINE SECRETIN-LIKE PROTEIN
16321. CYTOKINE SECRETIN-LIKE PROTEIN
16381. CYTOKINE SECRETIN-LIKE PROTEIN
16441. CYTOKINE SECRETIN-LIKE PROTEIN
16501. CYTOKINE SECRETIN-LIKE PROTEIN
16561. CYTOKINE SECRETIN-LIKE PROTEIN
16621. CYTOKINE SECRETIN-LIKE PROTEIN
16681. CYTOKINE SECRETIN-LIKE PROTEIN
16741. CYTOKINE SECRETIN-LIKE PROTEIN
16801. CYTOKINE SECRETIN-LIKE PROTEIN
16861. CYTOKINE SECRETIN-LIKE PROTEIN
16921. CYTOKINE SECRETIN-LIKE PROTEIN
16981. CYTOKINE SECRETIN-LIKE PROTEIN
17041. CYTOKINE SECRETIN-LIKE PROTEIN
17101. CYTOKINE SECRETIN-LIKE PROTEIN
17161. CYTOKINE SECRETIN-LIKE PROTEIN
17221. CYTOKINE SECRETIN-LIKE PROTEIN
17281. CYTOKINE SECRETIN-LIKE PROTEIN
17341. CYTOKINE SECRETIN-LIKE PROTEIN
17401. CYTOKINE SECRETIN-LIKE PROTEIN
17461. CYTOKINE SECRETIN-LIKE PROTEIN
17521. CYTOKINE SECRETIN-LIKE PROTEIN
17581. CYTOKINE SECRETIN-LIKE PROTEIN
17641. CYTOKINE SECRETIN-LIKE PROTEIN
17701. CYTOKINE SECRETIN-LIKE PROTEIN
17761. CYTOKINE SECRETIN-LIKE PROTEIN
17821. CYTOKINE SECRETIN-LIKE PROTEIN
17881. CYTOKINE SECRETIN-LIKE PROTEIN
17941. CYTOKINE SECRETIN-LIKE PROTEIN
18001. CYTOKINE SECRETIN-LIKE PROTEIN
18061. CYTOKINE SECRETIN-LIKE PROTEIN
18121. CYTOKINE SECRETIN-LIKE PROTEIN
18181. CYTOKINE SECRETIN-LIKE PROTEIN
18241. CYTOKINE SECRETIN-LIKE PROTEIN
18301. CYTOKINE SECRETIN-LIKE PROTEIN
18361. CYTOKINE SECRETIN-LIKE PROTEIN
18421. CYTOKINE SECRETIN-LIKE PROTEIN
18481. CYTOKINE SECRETIN-LIKE PROTEIN
18541. CYTOKINE SECRETIN-LIKE PROTEIN
18601. CYTOKINE SECRETIN-LIKE PROTEIN
18661. CYTOKINE SECRETIN-LIKE PROTEIN
18721. CYTOKINE SECRETIN-LIKE PROTEIN
18781. CYTOKINE SECRETIN-LIKE PROTEIN
18841. CYTOKINE SECRETIN-LIKE PROTEIN
18901. CYTOKINE SECRETIN-LIKE PROTEIN
18961. CYTOKINE SECRETIN-LIKE PROTEIN
19021. CYTOKINE SECRETIN-LIKE PROTEIN
19081. CYTOKINE SECRETIN-LIKE PROTEIN
19141. CYTOKINE SECRETIN-LIKE PROTEIN
19201. CYTOKINE SECRETIN-LIKE PROTEIN
19261. CYTOKINE SECRETIN-LIKE PROTEIN
19321. CYTOKINE SECRETIN-LIKE PROTEIN
19381. CYTOKINE SECRETIN-LIKE PROTEIN
19441. CYTOKINE SECRETIN-LIKE PROTEIN
19501. CYTOKINE SECRETIN-LIKE PROTEIN
19561. CYTOKINE SECRETIN-LIKE PROTEIN
19621. CYTOKINE SECRETIN-LIKE PROTEIN
19681. CYTOKINE SECRETIN-LIKE PROTEIN
19741. CYTOKINE SECRETIN-LIKE PROTEIN
19801. CYTOKINE SECRETIN-LIKE PROTEIN
19861. CYTOKINE SECRETIN-LIKE PROTEIN
19921. CYTOKINE SECRETIN-LIKE PROTEIN
19981. CYTOKINE SECRETIN-LIKE PROTEIN
20041. CYTOKINE SECRETIN-LIKE PROTEIN
20101. CYTOKINE SECRETIN-LIKE PROTEIN
20161. CYTOKINE SECRETIN-LIKE PROTEIN
20221. CYTOKINE SECRETIN-LIKE PROTEIN
20281. CYTOKINE SECRETIN-LIKE PROTEIN
20341. CYTOKINE SECRETIN-LIKE PROTEIN
20401. CYTOKINE SECRETIN-LIKE PROTEIN
20461. CYTOKINE SECRETIN-LIKE PROTEIN
20521. CYTOKINE SECRETIN-LIKE PROTEIN
20581. CYTOKINE SECRETIN-LIKE PROTEIN
20641. CYTOKINE SECRETIN-LIKE PROTEIN
20701. CYTOKINE SECRETIN-LIKE PROTEIN
20761. CYTOKINE SECRETIN-LIKE PROTEIN
20821. CYTOKINE SECRETIN-LIKE PROTEIN
20881. CYTOKINE SECRETIN-LIKE PROTEIN
20941. CYTOKINE SECRETIN-LIKE PROTEIN
21001. CYTOKINE SECRETIN-LIKE PROTEIN
21061. CYTOKINE SECRETIN-LIKE PROTEIN
21121. CYTOKINE SECRETIN-LIKE PROTEIN
21181. CYTOKINE SECRETIN-LIKE PROTEIN
21241. CYTOKINE SECRETIN-LIKE PROTEIN
21301. CYTOKINE SECRETIN-LIKE PROTEIN
21361. CYTOKINE SECRETIN-LIKE PROTEIN
21421. CYTOKINE SECRETIN-LIKE PROTEIN
21481. CYTOKINE SECRETIN-LIKE PROTEIN
21541. CYTOKINE SECRETIN-LIKE PROTEIN
21601. CYTOKINE SECRETIN-LIKE PROTEIN
21661. CYTOKINE SECRETIN-LIKE PROTEIN
21721. CYTOKINE SECRETIN-LIKE PROTEIN
21781. CYTOKINE SECRETIN-LIKE PROTEIN
21841. CYTOKINE SECRETIN-LIKE PROTEIN
21901. CYTOKINE SECRETIN-LIKE PROTEIN
21961. CYTOKINE SECRETIN-LIKE PROTEIN
22021. CYTOKINE SECRETIN-LIKE PROTEIN
22081. CYTOKINE SECRETIN-LIKE PROTEIN
22141. CYTOKINE SECRETIN-LIKE PROTEIN
22201. CYTOKINE SECRETIN-LIKE PROTEIN
22261. CYTOKINE SECRETIN-LIKE PROTEIN
22321. CYTOKINE SECRETIN-LIKE PROTEIN
22381. CYTOKINE SECRETIN-LIKE PROTEIN
22441. CYTOKINE SECRETIN-LIKE PROTEIN
22501. CYTOKINE SECRETIN-LIKE PROTEIN
22561. CYTOKINE SECRETIN-LIKE PROTEIN
22621. CYTOKINE SECRETIN-LIKE PROTEIN
22681. CYTOKINE SECRETIN-LIKE PROTEIN
22741. CYTOKINE SECRETIN-LIKE PROTEIN
22801. CYTOKINE SECRETIN-LIKE PROTEIN
22861. CYTOKINE SECRETIN-LIKE PROTEIN
22921. CYTOKINE SECRETIN-LIKE PROTEIN
22981. CYTOKINE SECRETIN-LIKE PROTEIN
23041. CYTOKINE SECRETIN-LIKE PROTEIN
23101. CYTOKINE SECRETIN-LIKE PROTEIN
23161. CYTOKINE SECRETIN-LIKE PROTEIN
23221. CYTOKINE SECRETIN-LIKE PROTEIN
23281. CYTOKINE SECRETIN-LIKE PROTEIN
23341. CYTOKINE SECRETIN-LIKE PROTEIN
23401. CYTOKINE SECRETIN-LIKE PROTEIN
23461. CYTOKINE SECRETIN-LIKE PROTEIN
23521. CYTOKINE SECRETIN-LIKE PROTEIN
23581. CYTOKINE SECRETIN-LIKE PROTEIN
23641. CYTOKINE SECRETIN-LIKE PROTEIN
23701. CYTOKINE SECRETIN-LIKE PROTEIN
23761. CYTOKINE SECRETIN-LIKE PROTEIN
23821. CYTOKINE SECRETIN-LIKE PROTEIN
23881. CYTOKINE SECRETIN-LIKE PROTEIN
23941. CYTOKINE SECRETIN-LIKE PROTEIN
24001. CYTOKINE SECRETIN-LIKE PROTEIN
24061. CYTOKINE SECRETIN-LIKE PROTEIN
24121. CYTOKINE SECRETIN-LIKE PROTEIN
24181. CYTOKINE SECRETIN-LIKE PROTEIN
24241. CYTOKINE SECRETIN-LIKE PROTEIN
24301. CYTOKINE SECRETIN-LIKE PROTEIN
24361. CYTOKINE SECRETIN-LIKE PROTEIN
24421. CYTOKINE SECRETIN-LIKE PROTEIN
24481. CYTOKINE SECRETIN-LIKE PROTEIN
24541. CYTOKINE SECRETIN-LIKE PROTEIN
24601. CYTOKINE SECRETIN-LIKE PROTEIN
24661. CYTOKINE SECRETIN-LIKE PROTEIN
24721. CYTOKINE SECRETIN-LIKE PROTEIN
24781. CYTOKINE SECRETIN-LIKE PROTEIN
24841. CYTOKINE SECRETIN-LIKE PROTEIN
24901. CYTOKINE SECRETIN-LIKE PROTEIN
24961. CYTOKINE SECRETIN-LIKE PROTEIN
25021. CYTOKINE SECRETIN-LIKE PROTEIN
25081. CYTOKINE SECRETIN-LIKE PROTEIN
25141. CYTOKINE SECRETIN-LIKE PROTEIN
25201. CYTOKINE SECRETIN-LIKE PROTEIN
25261. CYTOKINE SECRETIN-LIKE PROTEIN
25321. CYTOKINE SECRETIN-LIKE PROTEIN
25381. CYTOKINE SECRETIN-LIKE PROTEIN
25441. CYTOKINE SECRETIN-LIKE PROTEIN
25501. CYTOKINE SECRETIN-LIKE PROTEIN
25561. CYTOKINE SECRETIN-LIKE PROTEIN
25621. CYTOKINE SECRETIN-LIKE PROTEIN
25681. CYTOKINE SECRETIN-LIKE PROTEIN
25741. CYTOKINE SECRETIN-LIKE PROTEIN
25801. CYTOKINE SECRETIN-LIKE PROTEIN
25861. CYTOKINE SECRETIN-LIKE PROTEIN
25921. CYTOKINE SECRETIN-LIKE PROTEIN
25981. CYTOKINE SECRETIN-LIKE PROTEIN
26041. CYTOKINE SECRETIN-LIKE PROTEIN
26101. CYTOKINE SECRETIN-LIKE PROTEIN
26161. CYTOKINE SECRETIN-LIKE PROTEIN
26221. CYTOKINE SECRETIN-LIKE PROTEIN
26281. CYTOKINE SECRETIN-LIKE PROTEIN
26341. CYTOKINE SECRETIN-LIKE PROTEIN
26401. CYTOKINE SECRETIN-LIKE PROTEIN
26461. CYTOKINE SECRETIN-LIKE PROTEIN
26521. CYTOKINE SECRETIN-LIKE PROTEIN
26581. CYTOKINE SECRETIN-LIKE PROTEIN
26641. CYTOKINE SECRETIN-LIKE PROTEIN
26701. CYTOKINE SECRETIN-LIKE PROTEIN
26761. CYTOKINE SECRETIN-LIKE PROTEIN
26821. CYTOKINE SECRETIN-LIKE PROTEIN
26881. CYTOKINE SECRETIN-LIKE PROTEIN
26941. CYTOKINE SECRETIN-LIKE PROTEIN
27001. CYTOKINE SECRETIN-LIKE PROTEIN
27061. CYTOKINE SECRETIN-LIKE PROTEIN
27121. CYTOKINE SECRETIN-LIKE PROTEIN
27181. CYTOKINE SECRETIN-LIKE PROTEIN
27241. CYTOKINE SECRETIN-LIKE PROTEIN
27301. CYTOKINE SECRETIN-LIKE PROTEIN
27361. CYTOKINE SECRETIN-LIKE PROTEIN
27421. CYTOKINE SECRETIN-LIKE PROTEIN
27481. CYTOKINE SECRETIN-LIKE PROTEIN
27541. CYTOKINE SECRETIN-LIKE PROTEIN
27601. CYTOKINE SECRETIN-LIKE PROTEIN
27661. CYTOKINE SECRETIN-LIKE PROTEIN
27721. CYTOKINE SECRETIN-LIKE PROTEIN
27781. CYTOKINE SECRETIN-LIKE PROTEIN
27841. CYTOKINE SECRETIN-LIKE PROTEIN
27901. CYTOKINE SECRETIN-LIKE PROTEIN
27961. CYTOKINE SECRETIN-LIKE PROTEIN
28021. CYTOKINE SECRETIN-LIKE PROTEIN
28081. CYTOKINE SECRETIN-LIKE PROTEIN
28141. CYTOKINE SECRETIN-LIKE PROTEIN
28201. CYTOKINE SECRETIN-LIKE PROTEIN
28261. CYTOKINE SECRETIN-LIKE PROTEIN
28321. CYTOKINE SECRETIN-LIKE PROTEIN
28381. CYTOKINE SECRETIN-LIKE PROTEIN
28441. CYTOKINE SECRETIN-LIKE PROTEIN
28501. CYTOKINE SECRETIN-LIKE PROTEIN
28561. CYTOKINE SECRETIN-LIKE PROTEIN
28621. CYTOKINE SECRETIN-LIKE PROTEIN
28681. CYTOKINE SECRETIN-LIKE PROTEIN
28741. CYTOKINE SECRETIN-LIKE PROTEIN
28801. CYTOKINE SECRETIN-LIKE PROTEIN
28861. CYTOKINE SECRETIN-LIKE PROTEIN
289

1 AGACACCTCT GCCCTCACCA TGAGCCTCTG GCAGCCCCTG GTCCCTGGTGC TCCTGGTGCT
61 GGGCTGCTGC TTTGCTGCC CCAGACAGCG CCAGTCCACC CTTGTGCTCT TCCCTGGAGA
121 CCTGAGAAC AATCTCACCG ACAGGCAGCT GGCAAGAGGAA TACCTGTACC GCTATGGTTA
181 CACTCGGGTG GCAGAGATGC GTGGAGAGTC GAAATCTCTG GGGCCTGCAC TGCTGCTTCT
241 CCAGAACCAA CTGTCCCTGC CGCAGACCGG TGAGCTGGAT AGGCCACCGC TGAAGGCCAT
301 GCGAACCCCA CGGTGGGGGG TCCCAGACCT GGGCAGATTG CAAACCTTTG AGGGCGACCT
361 CAAGTGGCAC CACCACAACA TCACCTATTG GATCCAAAAC TACTCGGAAG ACTTGCCGCG
421 GGCGGTGATT GACGACGCC TTGCCCCGCCTT CGACGATGAC GAGTTGTGGT CCCTGGGCAA
481 CACCTTCACT CGCGTGTACA GCGGGGACGC AGACATCGTC ATCCAGTTTG GTGTCGGGA
541 GCACGGAGAC GGGTATCCCT TCGACGGGAA GGACGGGCTC CTGGCACACG CCTTTCCCTCC
601 TGGCCCCGGC ATTCAAGGGAG ACGCCCATT CGACGATGAC GAGTTGTGGT CCCTGGGCAA
661 GGGCGTCGTG GTTCCAACCTC GGTTTGGAAA CGCAGATGGC GCGGGCCTGCC ACTTCCCCCTT
721 CATCTTCAG GGGCGCTCCCT ACTCTGCTG CACCAACCGAC GGTGCGTCCG ACGGGTTGCC
781 CTGGTGCAGT ACCACGGCCA ACTACGACAC CGACGACCGG TTTGGCTTCT GCCCCAGCGA
841 GAGACTCTAC ACCCGGGACG GCAATGCTGA TGGGAAACCC TGCCAGTTTC CATTCACTT
901 CCAAGGCCAA TCCTACTCCG CCTGCACCCAC GGACGGTGC TCCGACGGCT ACCGCTGGTG
961 CGCCACCACC GCCAACATACG ACCGGGACAA GCTCTCGGC TTCTGCCCCGA CCCGAGCTGA
1021 CTGGACGGTG ATGGGGGCA ACTCGGGGGG GGAGCTGTGC GTCTTCCCCCT TCACTTTCCCT
1081 GGGTAAGGAG TACTCGACCT GTACCGACGA GGGCCGGGA GATGGGCGCC TCTGGTGC
1141 TACCACTCG AACTTTGACA GCGACAAGAA GTGGGGCTTC TGCCCGGACC AAGGATACAG
1201 TTTGTTCTC GTGGCGGC CGTACAGTTGG CCACGCGCTG GGCTTAGATC ATTCCCTCAGT
1261 GCCGGAGGCG CTCATGTACC CTATGTACCG CTTCACTGAG GGGCCCCCT TGCATAAGGA
1321 CGACGTGAAT GGCATCCGGC ACCCTATGG TCCTCGCCCT GAACCTGAGC CACGGCCTCC
1381 AACCAACCAC ACACCGCAGC CCACGGCTCC CCCGACGGTC TGCCCCACCG GACCCCCCAC
1441 TGTCCACCCCC TCAGAGCGCC CCACAGCTGG CCCCCACAGGT CCCCCCTCAG CTGGCCCCAC
1501 AGGTCCCCCCC ACTGCTGGCC CTTCTACGGC CACTACTGTG CCTTTGAGTC CGGTGGACGA
1561 TGCCCTGCAAC GTGAACATCT TCGACGCCAT CGCGGAGATT GGGAACCGAC TGTATTGTT
1621 CAAGGATGGG AAGTACTGGC GATTCTCTGA GGGCAGGGGG AGCCGGCCGC AGGGCCCCCTT
1681 CCTTATCGCC GACAAGTGGC CGCGCTGCC CCGCAAGCTG GACTCGGTCT TTGAGGAGCC
1741 GCTCTCCAAG AAGCTTTCT TCTTCTCTGG GCGCCAGGTG TGGGTGTACA CAGGCGCGTC
1801 GGTGCTGGGC CGAGGGCGTC TGACAAAGCT GGGCCTGGGA GCCGACGTGG CCCAGGTGAC
1861 CGGGGCCCTC CGGAGTGGCA GGGGGAAAGAT GCTGCTGTTG AGCGGGCGGC GCCTCTGGAG
1921 GTTCGACGTG AAGGCGCAGA TGGTGGATCC CGGGAGGCC AGCGAGGTGG ACCGGATGTT
1981 CCCCCGGGTG CCTTTGGACA CGCACGACGT CTTCAGTAC CGAGAGAAAG CCTATTCTG
2041 CCAGGACCGC TTCTACTGGC CGGTGAGTTG CCGGAGTGAG TTGAACCAAGG TGGACCAAGT
2101 GGGCTACGTG ACCTATGACA TCCTGCAGTG CCCTGAGGAC TAGGGCTCCC GTCCCTGCTT
2161 GCAGTGCCAT GTAAATCCCC ACTGGGACCA ACCCTGGGA AGGAGCCAGT TTGCCCCATA
2221 CAAACTGGTA TTCTGTTCTG GAGGAAAGGG AGGAGTGGAG GTGGGCTGGG CCCTCTCTTC
2281 TCACCTTGTGTTTGAGTGTCTA ATAAACTTGG ATTCTCTAAC CTTT

Figure 1

1

HUMAN OSTEOCLAST-SPECIFIC AND -RELATED GENES

RELATED APPLICATION

This application is a continuation of application Ser. No. 08/045,270 filed on Apr. 6, 1993 now abandoned.

BACKGROUND OF THE INVENTION

Excessive bone resorption by osteoclasts contributes to the pathology of many human diseases including arthritis, osteoporosis, periodontitis, and hypercalcemia of malignancy. During resorption, osteoclasts remove both the mineral and organic components of bone (Blair, H. C., et al., *J. Cell Biol.* 102:1164 (1986)). The mineral phase is solubilized by acidification of the sub-osteoclastic lacuna, thus allowing dissolution of hydroxyapatite (Vaes, G., *Clin. Orthop. Relat.* 231:239 (1988))). However, the mechanism(s) by which type I collagen, the major structural protein of bone, is degraded remains controversial. In addition, the regulation of osteoclastic activity is only partly understood. The lack of information concerning osteoclast function is due in part to the fact that these cells are extremely difficult to isolate as pure populations in large numbers. Furthermore, there are no osteoclastic cell lines available. An approach to studying osteoclast function that permits the identification of heretofore unknown osteoclast-specific or -related genes and gene products would allow identification of genes and gene products that are involved in the resorption of bone and in the regulation of osteoclastic activity. Therefore, identification of osteoclast-specific or -related genes or gene products would prove useful in developing therapeutic strategies for the treatment of disorders involving aberrant bone resorption.

10

15

20

25

30

35

SUMMARY OF THE INVENTION

The present invention relates to isolated DNA sequences encoding all or a portion of osteoclast-specific or -related gene products. The present invention further relates to DNA constructs capable of replicating DNA encoding osteoclast-specific or -related gene products. In another embodiment, the invention relates to a DNA construct capable of directing expression of all or a portion of the osteoclast-specific or -related gene product in a host cell.

40

Also encompassed by the present invention are prokaryotic or eukaryotic cells transformed or transfected with a DNA construct encoding all or a portion of an osteoclast-specific or -related gene product. According to a particular embodiment, these cells are capable of replicating the DNA construct comprising the DNA encoding the osteoclast-specific or -related gene product, and, optionally, are capable of expressing the osteoclast-specific or -related gene product. Also claimed are antibodies raised against osteoclast-specific or -related gene products, or portions of these gene products.

45

The present invention further embraces a method of identifying osteoclast-specific or -related DNA sequences and DNA sequences identified in this manner. In one embodiment, cDNA encoding osteoclast is identified as follows: First, human giant cell tumor of the bone was used to 1) construct a cDNA library; 2) produce ³²P-labelled cDNA to use as a stromal cell⁺, osteoclast⁻ probe, and 3) produce (by culturing) a stromal cell population lacking osteoclasts. The presence of osteoclasts in the giant cell tumor was confirmed by histological staining for the osteoclast marker, type 5 acid phosphatase (TRAP). In addition, monoclonal antibody reagents were used to characterize the multinucleated cells in the giant cell tumor, which cells were found to have a phenotype distinct from macrophages and consistent with osteoclasts.

50

55

60

65

2

clast marker, type 5 tartrate-resistant acid phosphatase (TRAP) and with the use of monoclonal antibody reagents.

The stromal cell population lacking osteoclasts was produced by dissociating cells of a giant cell tumor, then growing and passaging the cells in tissue culture until the cell population was homogeneous and appeared fibroblastic. The cultured stromal cell population did not contain osteoclasts. The cultured stromal cells were then used to produce a stromal cell⁺, osteoclast⁻ ³²P-labelled cDNA probe.

The cDNA library produced from the giant cell tumor of the bone was then screened in duplicate for hybridization to the cDNA probes: one screen was performed with the giant cell tumor cDNA probe (stromal cell⁺, osteoclast⁻), while a duplicate screen was performed using the cultured stromal cell cDNA probe (stromal cell⁺, osteoclast⁻). Hybridization to a stromal⁺, osteoclast⁺ probe, accompanied by failure to hybridize to a stromal⁺, osteoclast⁻ probe indicated that a clone contained nucleic acid sequences specifically expressed by osteoclasts.

In another embodiment, genomic DNA encoding osteoclast-specific or -related gene products is identified through known hybridization techniques or amplification techniques. In one embodiment, the present invention relates to a method of identifying DNA encoding an osteoclast-specific or -related protein, or gene product, by screening a cDNA library or a genomic DNA library with a DNA probe comprising one or more sequences selected from the group consisting of the DNA sequences set out in Table I (SEQ ID NOS: 1-32). Finally, the present invention relates to an osteoclast-specific or related protein encoded by a nucleotide sequence comprising a DNA sequence selected from the group consisting of the sequences set out in Table I, or their complementary strands.

BRIEF DESCRIPTION OF FIG. 1

The FIG. 1 shows cDNA sequence (SEQ ID NO: 33) of human gelatinase B, and highlights those portions of the sequence represented by the osteoclast-specific or -related cDNA clones of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

As described herein, Applicant has identified osteoclast-specific or osteoclast-related nucleic acid sequences. These sequences were identified as follows: Human giant cell tumor of the bone was used to 1) construct a cDNA library; 2) produce ³²P-labelled cDNA to use as a stromal cell⁺, osteoclast⁻ probe, and 3) produce (by culturing) a stromal cell population lacking osteoclasts. The presence of osteoclasts in the giant cell tumor was confirmed by histological staining for the osteoclast marker, type 5 acid phosphatase (TRAP). In addition, monoclonal antibody reagents were used to characterize the multinucleated cells in the giant cell tumor, which cells were found to have a phenotype distinct from macrophages and consistent with osteoclasts.

The stromal cell population lacking osteoclasts was produced by dissociating cells of a giant cell tumor, then growing the cells in tissue culture for at least five passages. After five passages the cultured cell population was homogeneous and appeared fibroblastic. The cultured population contained no multinucleated cells at this point, tested negative for type 5 acid phosphatase, and tested variably alkaline phosphatase positive. That is, the cultured stromal cell population did not contain osteoclasts. The cultured stromal

cells were then used to produce a stromal cell⁺, osteoclast⁻³²P-labelled cDNA probe.

The cDNA library produced from the giant cell tumor of the bone was then screened in duplicate for hybridization to the cDNA probes: one screen was performed with the giant cell tumor cDNA probe (stromal cell⁺, osteoclast⁻), while a duplicate screen was performed using the cultured stromal cell cDNA probe (stromal cell⁺ osteoclast⁻). Clones that hybridized to the giant cell tumor cDNA probe (stromal⁺, osteoclast⁻), but not to the stromal cell cDNA probe (stromal⁺, osteoclast⁻), were assumed to contain nucleic acid sequences specifically expressed by osteoclasts.

As a result of the differential screen described herein, DNA specifically expressed in osteoclast cells characterized as described herein was identified. This DNA, and equivalent DNA sequences, is referred to herein as osteoclast-specific or osteoclast-related DNA. Osteoclast-specific or -related DNA of the present invention can be obtained from sources in which it occurs in nature, can be produced recombinantly or synthesized chemically; it can be cDNA, genomic DNA, recombinantly-produced DNA or chemically-produced DNA. An equivalent DNA sequence is one which hybridizes, under standard hybridization conditions, to an osteoclast-specific or -related DNA identified as described herein or to a complement thereof.

Differential screening of a human osteoclastoma cDNA library was performed to identify genes specifically expressed in osteoclasts. Of 12,000 clones screened, 195 clones were identified which are either uniquely expressed in osteoclasts, or are osteoclast-related. These clones were further identified as osteoclast-specific, as evidenced by failure to hybridize to mRNA derived from a variety of unrelated human cell types, including epithelium, fibroblasts, lymphocytes, myelomonocytic cells, osteoblasts, and neuroblastoma cells. Of these, 32 clones contain novel cDNA sequences which were not found in the GenBank database.

A large number of cDNA clones obtained by this procedure were found to represent 92 kDa type IV collagenase (gelatinase B; E.C. 3.4.24.35) as well as tartrate resistant acid phosphatase. In situ hybridization localized mRNA for gelatinase B to multinucleated giant cells in human osteoclastomas. Gelatinase B immunoreactivity was demonstrated in giant cells from 8/8 osteoclastomas, osteoclasts in normal bone, and in osteoclasts of Paget's disease by use of a polyclonal antisera raised against a synthetic gelatinase B peptide. In contrast, no immunoreactivity for 72 kDa type IV collagenase (gelatinase A; E.C. 3.4.24.24), which is the product of a separate gene, was detected in osteoclastomas or normal osteoclasts.

The present invention has utility for the production and identification of nucleic acid probes useful for identifying osteoclast-specific or -related DNA. Osteoclast-specific or -related DNA of the present invention can be used to produce osteoclast-specific or -related gene products useful in the therapeutic treatment of disorders involving aberrant bone resorption. The osteoclast-specific or -related sequences are also useful for generating peptides which can then be used to produce antibodies useful for identifying osteoclast-specific or -related gene products, or for altering the activity of osteoclast-specific or -related gene products. Such antibodies are referred to as osteoclast-specific antibodies. Osteoclast-specific antibodies are also useful for identifying osteoclasts. Finally, osteoclast-specific or -related DNA sequences of the present invention are useful in gene therapy. For example, they can be used to alter the

expression in osteoclasts of an aberrant osteoclast -specific or -related gene product or to correct aberrant expression of an osteoclast-specific or -related gene product. The sequences described herein can further be used to cause osteoclast-specific or related gene expression in cells in which such expression does not ordinarily occur, i.e., in cells which are not osteoclasts.

Example 1—Osteoclast cDNA Library Construction

Messenger RNA (mRNA) obtained from a human osteoclastoma ('giant cell tumor of bone'), was used to construct an osteoclastoma cDNA library. Osteoclastomas are actively bone resorptive tumors, but are usually non-metastatic. In cryostat sections, osteoclastomas consist of ~30% multinucleated cells positive for tartrate resistant acid phosphatase (TRAP), a widely utilized phenotypic marker specific in vivo for osteoclasts (Minkin, *Calcif. Tissue Int.* 34:285-290 (1982)). The remaining cells are uncharacterized 'stromal' cells, a mixture of cell types with fibroblastic/mesenchymal morphology. Although it has not yet been definitively shown, it is generally held that the osteoclasts in these tumors are non-transformed, and are activated to resorb bone in vivo by substance(s) produced by the stromal cell element.

Monoclonal antibody reagents were used to partially characterize the surface phenotype of the multinucleated cells in the giant cell tumors of long bone. In frozen sections, all multinucleated cells expressed CD68, which has previously been reported to define an antigen specific for both osteoclasts and macrophages (Horton, M. A. and M. H. Helfrich, In *Biology and Physiology of the Osteoclast*, B. R. Rifkin and C. V. Gay, editors, CRC Press, Inc. Boca Raton, Fla., 33-54 (1992)). In contrast, no staining of giant cells was observed for CD11b or CD14 surface antigens, which are present on monocyte/macrophages and granulocytes (Arnaout, M. A. et al. *J. Cell. Physiol.* 137:305 (1988); Haziot, A. et al. *J. Immunol.* 141:547 (1988)). Cytocentrifuge preparations of human peripheral blood monocytes were positive for CD68, CD11b, and CD14. These results demonstrate that the multinucleated giant cells of osteoclastomas have a phenotype which is distinct from that of macrophages, and which is consistent with that of osteoclasts.

Osteoclastoma tissue was snap frozen in liquid nitrogen and used to prepare poly A⁺ mRNA according to standard methods. cDNA cloning into a pcDNAII vector was carried out using a commercially-available kit (Librarian, InVitrogen). Approximately 2.6x10⁶ clones were obtained, >95% of which contained inserts of an average length 0.6 kB.

Example 2—Stromal Cell mRNA Preparation

A portion of each osteoclastoma was snap frozen in liquid nitrogen for mRNA preparation. The remainder of the tumor was dissociated using brief trypsinization and mechanical disaggregation, and placed into tissue culture. These cells were expanded in Dulbecco's MEM (high glucose, Sigma) supplemented with 10% newborn calf serum (MA Bioproducts), gentamycin (0.5 mg/ml), L-glutamine (2 mM) and non-essential amino acids (0.1 mM) (Gibco). The stromal cell population was passaged at least five times, after which it showed a homogenous, fibroblastic looking cell population that contained no multinucleated cells. The stromal cells were mononuclear, tested negative acid phosphatase, and tested variably alkaline phosphatase positive. These findings indicate that propagated stromal cells (i.e., stromal cells that

are passaged in culture) are non-osteoclastic and non-activated.

Example 3—Identification of DNA Encoding Osteoclastoma-Specific or -Related Gene Products by Differential Screening of an Osteoclastoma cDNA Library

A total of 12,000 clones drawn from the osteoclastoma cDNA library were screened by differential hybridization, using mixed ^{32}P labelled cDNA probes derived from (1) giant cell tumor mRNA (stromal cell $^+$, OC $^+$), and (2) mRNA from stromal cells (stromal cell $^+$, OC $^-$) cultivated from the same tumor. The probes were labelled with ^{32}P dCTP by random priming to an activity of $\sim 10^9 \text{ CPM}/\mu\text{g}$. Of these 12,000 clones, 195 gave a positive hybridization signal with giant cell (i.e., osteoclast and stromal cell) mRNA, but not with stromal cell mRNA. Additionally, these clones failed to hybridize to cDNA produced from mRNA derived from a variety of unrelated human cell types including epithelial cells, fibroblasts, lymphocytes, myelomonocytic cells, osteoblasts, and neuroblastoma cells. The failure of these clones to hybridize to cDNA produced from mRNA derived from other cell types supports the conclusion that these clones are either uniquely expressed in osteoclasts, or are osteoclast-related.

The osteoclast (OC) cDNA library was screened for differential hybridization to OC cDNA (stromal cell $^+$, OC $^+$) and stromal cell cDNA (stromal cell $^+$, OC $^-$) as follows:

NYTRAN filters (Schleicher & Schuell) were placed on agar plates containing growth medium and ampicillin. Individual bacterial colonies from the OC library were randomly picked and transferred, in triplicate, onto filters with pre-ulcd grids and then onto a master agar plate. Up to 200 colonies were inoculated onto a single 90-mm filter/plate using these techniques. The plates were inverted and incubated at 37° C. until the bacterial inoculates had grown (on the filter) to a diameter of 0.5–1.0 mm.

The colonies were then lysed, and the DNA bound to the filters by first placing the filters on top of two pieces of Whatman 3 MM paper saturated with 0.5N NaOH for 5 minutes. The filters were neutralized by placing on two pieces of Whatman 3 MM paper saturated with 1M Tris-HCL, pH 8.0 for 3–5 minutes. Neutralization was followed by incubation on another set of Whatman 3 MM papers saturated with 1M Tris-HCL, pH 8.0/1.5M NaCl for 3–5 minutes. The filters were then washed briefly in 2xSSC.

DNA was immobilized on the filters by baking the filters at 80° C. for 30 minutes. Filters were best used immediately, but they could be stored for up to one week in a vacuum jar at room temperature.

Filters were prehybridized in 5–8 ml of hybridization solution per filter, for 2–4 hours in a heat sealable bag. An additional 2 ml of solution was added for each additional filter added to the hybridization bag. The hybridization

buffer consisted of 5xSSC, 5xDenhardt's solution, 1% SDS and 100 µg/ml denatured heterologous DNA.

Prior to hybridization, labeled probe was denatured by heating in 1xSSC for 5 minutes at 100° C., then immediately chilled on ice. Denatured probe was added to the filters in hybridization solution, and the filters hybridized with continuous agitation for 12–20 hours at 65° C.

After hybridization, the filters were washed in 2xSSC/0.2% SDS at 50°–60° C. for 30 minutes, followed by washing in 0.2xSSC/0.2% SDS at 60° C. for 60 minutes.

The filters were then air dried and autoradiographed using an intensifying screen at -70° C. overnight.

Example 4—DNA Sequencing of Selected Clones

Clones reactive with the mixed tumor probe, but unreactive with the stromal cell probe, are expected to contain either osteoclast-related, or *in vivo* 'activated' stromal-cell-related gene products. One hundred and forty-four cDNA clones that hybridized to tumor cell cDNA, but not to stromal cell cDNA, were sequenced by the dideoxy chain termination method of Sanger et al. (Sanger F, et al. *Proc. Natl. Acad. Sci. USA* 74:5463 (1977)) using sequenase (US Biochemical). The DNASIS (Hitatchi) program was used to carry out sequence analysis and a homology search in the GenBank/EMBL database.

Fourteen of the 195 tumor $^+$ stromal $^-$ clones were identified as containing inserts with a sequence identical to the osteoclast marker, type 5 tartrate-resistant acid phosphatase (TRAP) (GenBank accession number J04430 M19534). The high representation of TRAP positive clones also indicates the effectiveness of the screening procedure in enriching for clones which contain osteoclast-specific or related cDNA sequences.

Interestingly, an even larger proportion of the tumor $^+$ stromal $^-$ clones (77/195; 39.5%) were identified as human gelatinase B (macrophage-derived gelatinase) (Wilhelm, S. M. *J. Biol. Chem.* 264:17213 (1989)), again indicating high expression of this enzyme by osteoclasts. Twenty-five of the gelatinase B clones were identified by dideoxy sequence analysis; all 25 showed 100% sequence homology to the published gelatinase B sequence (Genbank accession number J05070). The portions of the gelatinase B cDNA sequence covered by these clones is shown in the FIGURE (SEQ ID NO: 33). An additional 52 gelatinase B clones were identified by reactivity with a ^{32}P -labelled probe for gelatinase B.

Thirteen of the sequenced clones yielded no readable sequence. A DNASIS search of GenBank/EMBL databases revealed that, of the remaining 91 clones, 32 clones contain novel sequences which have not yet been reported in the databases or in the literature. These partial sequences are presented in Table I. Note that three of these sequences were repeats, indicating fairly frequent representation of mRNA related to this sequence. The repeat sequences are indicated by^{a, b} superscripts (Clones 198B, 223B and 32C of Table I).

TABLE I

PARTIAL SEQUENCES OF 32 NOVEL OC-SPECIFIC OR -RELATED EXPRESSED GENES (cDNA CLONES)

34A (SEQ ID NO: 1)					
1 GCAAATATCT	AAGTTTATTG	CTTGGATTTC	TAGTGAGAGC	TGTGAAATT	GGTGATGTCA
61 AATGTTCTA	GGGTTTTTTT	AGTTTGTTT	TATTGAAAAA	TTTAATTAT	TATGCTATAG
121 GTGATATCT	CTTTGAATAA	ACCTATAATA	GAAAATAGCA	GCAGACAACA	
4B (SEQ ID NO: 2)					
1 GTGTCAACCT	GCATATCCCA	AAAATGTCAA	AATGCTGCAT	CTGGTTAACG	TCCGGGTAGG

TABLE I-continued

PARTIAL SEQUENCES OF 32 NOVEL OC-SPECIFIC OR -RELATED EXPRESSED GENES (cDNA CLONES)

61	GGG					
12B	(SEQ ID NO: 3)					
1	CTTCCCCTCTC	TTCCTTCCCT	TTCCCAAGCA	GAGGTGCTCA	CTCCATGGCC	ACCGCCACCA
61	CAGGCCACCA	CGGAGTACTG	CCAGACTACT	GCTGATGTTG	TCTTAAGGCC	CAGGGAGTCT
121	CAACCAGCTG	GTGGTGAATG	CTGGCTGGCA	CGGGACCCCC	CCC	
28B	(SEQ ID NO: 4)					
1	TTTATTTGT	AAATATATGT	ATTACATCCC	TAGAAAAAGA	ATCCAGGAT	TTTCCCTCT
61	GTGTGTTTC	GTCTTGCTTC	TTCATGGTCC	ATGATGCCAG	CTGAGGTTGT	CAGTACAATG
121	AAACCAAAT	GGCGGGTGG	AAGCAGATTA	TTCTGCCATT	TTTCAGGTC	TTT
37B	(SEQ ID NO: 5)					
1	GGCTGGACAT	GGGTGCCCTC	CACGTCCCTC	ATATCCCCAG	GCACACTCTG	GCCTCAGGTT
61	TTGGCCCTGGC	CATGTCATCT	ACCTGGAGTG	GGCCCTCCCC	TTCTTCAGCC	TTGAAATCAA
121	AGCCACTTTG	TTAGGGAGG	ATTTCCAGA	CCACTCATCA	CATAAAAAAA	TATTTGAAA
181	ACAAAAAAA	AAAAAAA				
55B	(SEQ ID NO: 6)					
1	TTGACAAAGC	TGTTTATTC	CACCAATAAA	TAGTATATGG	TGATTGGGT	TTCTATTTAT
61	AAGAGTAGTG	GCTTATTATAT	GGGGTATCAT	TTGATGCTC	ATAAATAGT	CATATCTACT
121	TAATTGCTT	TC				
60B	(SEQ ID NO: 7)					
1	GAAGAGAGTT	GTATGTACAA	CCCCAACAGG	CAAGGCAGCT	AAATGCAGAG	GGTACAGAGA
61	GATCCCGAGG	GAATT				
86B	(SEQ ID NO: 8)					
1	GGATGAAAC	ATGAGAAGT	CCAGAGAAAA	ACAATTAA	AAAAGGTGG	AAAAGTTACG
61	GCAAACCTGA	GATTTCAGCA	TAACATCTT	AGTTAGAAGT	GAAGAAAAGA	AGAGGGAGGC
121	TGGTGGCTGT	TGCACTATC	AAATAGTTAT	C		
87B	(SEQ ID NO: 9)					
1	TTCTTGATCT	TTAGAACACT	ATGAATAGGG	AAAAAAGAAA	AAACTGTICA	AAATAAAATG
61	TAGGAGCCGT	GCTTTGGA	TGCTTGAGTG	AGGAGCTCAA	CAAGCTCT	CCCAAGAAAG
181	CAATGATAAA	ACTTGACAAA	A			
98B	(SEQ ID NO: 10)					
1	ACCAATTCT	AACAATTTT	ACTGAAAAAT	TTTTGGTCAA	AGTTCTAAGC	TTAATCACAT
61	CTAAAGAAT	AGAGGAATA	TATAGCCCAT	CTTACTAGAC	ATACAGTATT	AAACTGGACT
121	GAATATGAGG	ACAACCTCA	GTGGTCATTA	AACCCCTCA	AA	
110B	(SEQ ID NO: 11)					
1	ACATATATTA	ACAGCATTCA	TTTGGCCAAA	ATCTACACGT	TTGAGAAC	CTACTGTATA
61	TAAGATGGGA	ATGATATCAAG	TATAGACTAT	GAAGATGCAA	ATAACAAGTC	AAGGTTAGAT
121	TAACCTTTT	TTTTACATT	ATAAAAATTAA	CTTGT		
118B	(SEQ ID NO: 12)					
1	CCAAATTCT	CTGGAAATCCA	TCCTCCCTCC	CATCACCATA	GCCTCGAGAC	GTCATTTCTG
61	TTTGACTACT	CCAGC				
133B	(SEQ ID NO: 13)					
1	AACTAACCTC	CTGGGACCCC	TGCCTCACTC	ATTACACCA	ACCAACCAAC	TATCTATAAA
61	CCTGAGCCAT	GGCCATCCCT	TATGAGCGGC	GCAGTGATTA	TAGGCTTTCG	CTCTAAGATA
121	AAAT					
140B	(SEQ ID NO: 14)					
1	ATTATTATTC	TTTTTTATG	TTAGCTTAGC	CATGCAAAAT	TTACTGGTGA	AGCAGTTAAAT
61	AAAACACACA	TCCCATTGAA	GGGTTTTGTA	CATTTCAGTC	CTTACAATA	ACAAAGCAAT
121	GATAAACCG	GCACGTCTG	ATAGGAAATT	C		
144B	(SEQ ID NO: 15)					
1	CGTGACACAA	ACATGCACTC	TTTTTATTCA	AAAAACAGCC	TOGTTTCCTA	AAACAATACA
61	AACAGCATGT	TCATCAGCA	GAAGCTGGCC	GTGGGCAGGG	GGGCC	
198B*	(SEQ ID NO: 16)					
1	ATAGGTTAGA	TCTCATCA	CGGGACTAGT	TAGCTTAAAG	CACCTAGAG	GACTAGGCTA
61	ATCTGACTTC	TCACTTCTA	ACTTCCCTCT	TATATCCCA	GTCTATGTT	
121	TCTACTCCAA	TTTCATAAATC	TATTCTAAAG	TCCTTGGTAC	AAAGTACATG	ATAAAAAGAA
181	ATGTGATTG	TCTTCCCTTC	TTTGACTTT	TRAATAAAAG	TATTTATCTC	CTGTCACAG
241	TTTAAT					
212B	(SEQ ID NO: 17)					
1	GTCCAGTATA	AAAGAAAAGCG	TIAAATGCGGT	AAGCTAGAGG	ATGTAATA	TCTTTTATGT
61	CCTCTAGATA	AAACACCCGA	TTAACAGATG	TTAACCTTT	ATGTTTTGAT	TTGCTTAAAG
121	AATGCCCTC	TACACATTAG	CTCCAGCTAA	AAAGACACAT	TGAGAGCTTA	GAGGATAGTC
181	TCTGAGC					
223B*	(SEQ ID NO: 18)					
1	GCACTTGGAA	GGGAGTTGGT	GTGCTATTIT	TGAAGCAGAT	GTGGTGATAC	TGAGATTGTC
61	TGTCAGTTT	CCCCATTGAT	TTGTCCTCA	AATGATCCTT	CCTACTTTGC	TTCTCTCCAC
121	CCATGACCTT	TTTCACITG	GCCATCAAGG	ACTTCCCTGA	CAGCTTGTGT	ACTCTTAGGC
181	TAAGAGATGT	GAATCAGCC	TGCCCCGTAC	TG		
241B	(SEQ ID NO: 19)					
1	TGTTAGTTT	TAGGAAGGCC	TGTCTCTGG	GAGTGAGGTT	TATTAGTCGA	CTTCCTGGAG
61	CTAGACGTCC	TATAGTTAGT	CACTGGGGAT	GGTGAAAGAG	GGAGAAGAGG	AAGGGCGAAG
121	GGAAAGGGCTC	TTTGCTAGTA	TCTCCATTTC	TAGAAGATGG	TTTAGATGAT	AACACAGGT
181	CTATATGAGC	ATAGTAAGGC	TGT			
32C*	(SEQ ID NO: 20)					
1	CCTATTCTCG	ATCCCTGACTT	TGGACAAGGC	CCTTCAGCCA	GAAGACTGAC	AAAGTCATCC
121	TCCGTCTACC	AGAGCGTGC	CTTGTGATCC	TAATAAAGC	TTCATCTCG	GCTGTGCCCT
161	GGGTGGAAAGG	CGCAGGATTC	TGCAGCTGCT	TTTCATTC	TCTTCCTAA	TTTCATT

TABLE I-continued

PARTIAL SEQUENCES OF 32 NOVEL OC-SPECIFIC OR -RELATED EXPRESSED GENES (cDNA CLONES)

34C (SEQ ID NO: 21)					
1 CGGAGCCGTAG	GTTGTTTAT	TCTTGACAA	ATCATTACAA	AACCAAGTCT	GGGGCAGTCA
61 CGGCCCCCAC	CCATCACCCC	AGTGAATGG	CTAGCTGCTG	GCCTTT	
47C (SEQ ID NO: 22)					
1 TTAGTCAGT	CAAAGCAGGC	AACCCCCCTT	GGCACTGCTG	CCACTGGGGT	CATGGCGGTT
61 GTGGCAGCTG	GGAGGGTTTC	CCCAACACCC	TCCTCTGCTT	CCCTGTGTG	CGGGGTCTCA
121 GGAGCTGACC	CAGACTGGA				
65C (SEQ ID NO: 23)					
1 GCTGAAGTGT	TAAGAGAGAT	TTTGGCTTA	AAGGCTTCAT	CATGAAAGTG	TACATGCATA
61 TGCAAGTGTG	AAITACGTGG	TATGGATGGT	TGCTTGTGTTA	TTAACATAAG	ATGACAGCA
121 AACTGCCGT	TTAGACTCTT	CTTAATATG	ATGTOCTAAC	ACTGGGCTG	CTTATGC
79C (SEQ ID NO: 24)					
1 GGCACTGGGA	TATGGAATCC	AGAAGGGAAA	CAAGCACTGG	ATAATTTAAA	ACAGCTGGGG
61 AGAAAACCTGG	GGAAACAAAG	GATAATATCT	CATGGCTCGA	ATAAGAAACA	AOGCCTGTGG
121 CATTGCAAC	CTGGCCAGCT	TOCCCAGAT	GTGACTCCAG	CCAGAAA	
84C (SEQ ID NO: 25)					
1 GCCAGGGGG	ACCGTCCTTA	TTCCCTCTCT	GCCTCAGAGG	TCAGGAAGGA	GGTCTGGCAG
61 GACCTGCACT	GGGCCCCTAGT	CATCTGTGGC	AGCGAAGGTG	AAGGGACTCA	CCTTGTGCGC
121 CGTGCCGTGAG	TAGAACTTGT	TCTGGAATTC	C		
86C (SEQ ID NO: 26)					
1 AACCTTTICA	CACTCTGGTA	TTTTTAGTT	AACAATATAT	GTGTTGTGTC	TTGAAAATA
61 GTTCATATCA	ATTCTATATG	AGCTGTCTCA	TTCTTTTTT	ATAGTCATA	TACAGTAGTA
121 TTCAATTATA	AGAATATATC	CTAATACCTT	TTAAAAA		
87C (SEQ ID NO: 27)					
1 GGATAAGAAA	GAAGGCCCTGA	GGCCTAGGGG	CCGRGGCTGG	CCTGGCTCTC	AGTOCTGGGA
61 CGCAGCAGCC	CGCACAGCTT	GAGAGGGGCA	CTTCCTCTTG	CTTAGGTTGG	TGAGGATCTG
121 GTCCTGGTTG	GGCCGGTGGAG	AGGCCACAAA			
88C (SEQ ID NO: 28)					
1 CTGACCTTCG	AGAGTTTGAC	CTGGAGCCGG	ATACCTACTG	CCGCTATGAC	TCGGTCAGCG
61 TGTCAACCG	AGCCGTGAGC	GACGACTCCG	GTGGGAAAGT	TCTGCGGGCA	T
89C (SEQ ID NO: 29)					
1 ATCCCTGGCT	GTGGATAGTC	CTTTTGTGTA	GCAAATGCTC	OCTCTTAAAG	GTATAGGGC
61 TCCCTGAGTT	TGGGAGTGTG	GAAGTACTAC	TTAACGTCT	GTCTGCTTG	GCTGTOGTA
121 TCGTTTCTG	GTGATGTTGT	GCTAACATA	AGAATAC		
101C (SEQ ID NO: 30)					
1 GGCTGGCCAT	CCCTCTCTC	CTCCATCCCC	ATACATCACC	AGGTCTAAATG	TTTACAAACG
61 GTGCCAGCCC	GGCTCTGAAG	CCAAGGGCCG	TCCGTGCCAC	GGTGGCTGTG	AGTATTCCTC
121 CGTTAGCTT	CCCATAAAGT	TGGAGTACT	GC		
112C (SEQ ID NO: 31)					
1 CCAACTCTTA	CCGGCATAACA	GACCCACAGA	GTGCCATCCC	TGAGAGACCA	GACCGCTCCC
161 CAATACTCTC	CTAAAAATAAA	CATGAAGCAC			
114C (SEQ ID NO: 32)					
1 CATGGATGAA	TGTCTCATGG	TGGGAAGGAA	CATGGTACAT	TTC	

^aRepeated 3 times^bRepeated 2 times

Sequence analysis of the OC⁺ stromal cell⁻ cloned DNA sequences revealed, in addition to the novel sequences, a number of previously-described genes. The known genes identified (including type 5 acid phosphatase, gelatinase B, cystatin C (13 clones), Alu repeat sequences (11 clones), creatine kinase (6 clones) and others) are summarized in Table II. In situ hybridization (described below) directly demonstrated that gelatinase B mRNA is expressed in multi-nucleated osteoclasts and not in stromal cells. Although gelatinase B is a well-characterized protease, its expression at high levels in osteoclasts has not been previously described. The expression in osteoclasts of cystatin C, a cysteine protease inhibitor, is also unexpected. This finding has not yet been confirmed by in situ hybridization. Taken together, these results demonstrate that most of these identified genes are osteoclast-expressed, thereby confirming the effectiveness of the differential screening strategy for identifying DNA encoding osteoclast-specific or -related gene products. Therefore, novel genes identified by this method have a high probability of being OC-specific or related.

In addition, a minority of the genes identified by this screen are probably not expressed by OCs (Table II). For example, type III collagen (6 clones), collagen type I (1 clone), dermatansulfate (1 clone), and type VI collagen (1

clone) are more likely to originate from the stromal cells or from osteoblastic cells which are present in the tumor. These cDNA sequences survive the differential screening process either because the cells which produce them in the tumor in vivo die out during the stromal cell propagation phase, or because they stop producing their product in vitro. These clones do not constitute more than 5-10% of the all sequences selected by differential hybridization.

TABLE II

SEQUENCE ANALYSIS OF CLONES ENCODING KNOWN SEQUENCES FROM AN OSTEOCLASTOMA cDNA LIBRARY

Clones with Sequence Homology to Collagenase Type IV	25 total
Clones with Sequence Homology to Type 5 Taurate Resistant Acid Phosphatase	14 total
Clones with Sequence Homology to Cystatin C	13 total
Clones with Sequence Homology to Alu-repeat Sequences	11 total
Clones with Sequence Homology to Creatine Kinase	6 total
Clones with Sequence Homology to	6 total

TABLE II-continued

SEQUENCE ANALYSIS OF CLONES ENCODING KNOWN SEQUENCES FROM AN OSTEOCLASTOMA cDNA LIBRARY		
Type III Collagen	5 total	
Clones with Sequence Homology to MHC Class I γ Invariant Chain	3 total	
Clones with Sequence Homology to MHC Class II β Chain	10 total	
One or Two Clone(s) with Sequence Homology to Each of the Following:		
α 1 collagen type I		
γ interferon inducible protein		
osteoponin		
Human chondroitin/dermatan sulfate	15	
α globin		
β glucuronidase/sphingolipid activator		
Human CAPL protein (Ca binding)		
Human EST 01024		
Type VI collagen		
Human EST 00553		

Example 5—In situ Hybridization of OC-Expressed Genes

In situ hybridization was performed using probes derived from novel cloned sequences in order to determine whether the novel putative OC-specific or -related genes are differentially expressed in osteoclasts (and not expressed in the stromal cells) of human giant cell tumors. Initially, in situ hybridization was performed using antisense (positive) and sense (negative control) cRNA probes against human type IV collagenase/gelatinase B labelled with 35 S-UTP.

A thin section of human giant cell tumor reacted with the antisense probe resulted in intense labelling of all OCs, as indicated by the deposition of silver grains over these cells, but failed to label the stromal cell elements. In contrast, only minimal background labelling was observed with the sense (negative control) probe. This result confirmed that gelatinase B is expressed in human OCs.

In situ hybridization was then carried out using cRNA probes derived from 11/32 novel genes, labelled with digoxigenin UTP according to known methods.

The results of this analysis are summarized in Table III. Clones 28B, 118B, 140B, 198B, and 212B all gave positive reactions with OCs in frozen sections of a giant cell tumor, as did the positive control gelatinase B. These novel clones therefore are expressed in OCs and fulfill all criteria for OC-relatedness. 198B is repeated three times, indicating relatively high expression. Clones 4B, 37B, 88C and 98B produced positive reactions with the tumor tissue; however the signal was not well-localized to OCs. These clones are therefore not likely to be useful and are eliminated from further consideration. Clones 86B and 87B failed to give a positive reaction with any cell type, possibly indicating very low level expression. This group of clones could still be useful but may be difficult to study further. The results of this analysis show that 5/11 novel genes are expressed in OCs, indicating that ~50% of novel sequences likely to be OC-related.

To generate probes for the in situ hybridizations, cDNA derived from novel cloned osteoclast-specific or -related cDNA was subcloned into a BlueScript II SK(-) vector. The orientation of cloned inserts was determined by restriction analysis of subclones. The T7 and T3 promoters in the BlueScriptII vector was used to generate 35 S-labelled (35 S-UTP 850 Ci/mmol, Amersham, Arlington Heights, Ill.), or

12
UTP digoxigenin labelled cRNA probes.

TABLE III

Clone	Reactivity with:	
	Osteoclasts	Stromal Cells
4B	+	+
28B*	+	-
37B	+	+
86B	-	-
87B	-	-
88C	+	+
98B	+	+
118B*	+	-
140B*	+	-
198B*	+	-
212B*	+	-
Gelatinase B*	+	-

*OC-expressed, as indicated by reactivity with antisense probe and lack of reactivity with sense probe on OCs only.

In situ hybridization was carried out on 7 micron cryostat sections of a human osteoclastoma as described previously (Chang, L.-C. et al. *Cancer Res.* 49:6700 (1989)). Briefly, tissue was fixed in 4% paraformaldehyde and embedded in OCT (Miles Inc., Kankakee, Ill.). The sections were rehydrated, postfixed in 4% paraformaldehyde, washed, and pretreated with 10 mM DTT, 10 mM iodoacetamide, 10 mM N-ethylmaleimide and 0.1 triethanolamine-HCl. Prehybridization was done with 50% deionized formamide, 10 mM Tris-HCl, pH 7.0, 1x Denhardt's, 500 mg/ml tRNA, 80 mg/ml salmon sperm DNA, 0.3M NaCl, mM EDTA, and 100 mM DTT at 45° C. for 2 hours. Fresh hybridization solution containing 10% dextran sulfate and 1.5 ng/ml 35 S-labelled or digoxigenin labelled RNA probe was applied after heat denaturation. Sections were coverslipped and then incubated in a moistened chamber at 45°–50° C. overnight. Hybridized sections were washed four times with 50% formamide, 2x SSC, containing 10 mM DTT and 0.5% Triton X-100 at 45° C. Sections were treated with RNase A and RNase T1 to digest single-stranded RNA, washed four times in 2x SSC/10 mM DTT.

In order to detect 35 S-labelling by autoradiography, slides were dehydrated, dried, and coated with Kodak NTB-2 emulsion. The duplicate slides were split, and each set was placed in a black box with desiccant, sealed, and incubated at 4° C. for 2 days. The slides were developed (4 minutes) and fixed (5 minutes) using Kodak developer D19 and Kodak fixer. Hematoxylin and eosin were used as counterstains.

In order to detect digoxigenin-labelled probes, a Nucleic Acid Detection Kit (Boehringer-Mannheim, Cat #1175041) was used. Slides were washed in Buffer 1 consisting of 100 mM Tris/150 mM NaCl, pH 7.5, for 1 minute. 100 μ l Buffer 2 was added (made by adding 2 mg/ml blocking reagent as provided by the manufacturer) in Buffer 1 to each slide. The slides were placed on a shaker and gently swirled at 20° C.

Antibody solutions were diluted 1:100 with Buffer 2 (as provided by the manufacturer). 100 μ l of diluted antibody solution was applied to the slides and the slides were then incubated in a chamber for 1 hour at room temperature. The slides were monitored to avoid drying. After incubation with antibody solution, slides were washed in Buffer 1 for 10 minutes, then washed in Buffer 3 containing 2 mM levamisole for 2 minutes.

After washing, 100 μ l color solution was added to the slides. Color solution consisted of nitroblue/tetrazolium salt

13

(NBT) (1:225 dilution) 4.5 μ l, 5-bromo-4-chloro-3-indolyl phosphate (1:285 dilution) 3.5 μ l, levamisole 0.2 mg in Buffer 3 (as provided by the manufacturer) in a total volume of 1 ml. Color solution was prepared immediately before use.

After adding the color solution, the slides were placed in a dark, humidified chamber at 20° C. for 2-5 hours and monitored for color development. The color reaction was stopped by rinsing slides in TE Buffer.

The slides were stained for 60 seconds in 0.25% methyl green, washed with tap water, then mounted with water-based Permount (Fisher).

Example 6—Immunohistochemistry

15 Immunohistochemical staining was performed on frozen and paraffin embedded tissues as well as on cytopsin preparations (see Table IV). The following antibodies were used: polyclonal rabbit anti-human gelatinase antibodies; Ab110 for gelatinase B; monoclonal mouse anti-human CD68 antibody (clone KP1) (DAKO, Denmark); Mol (anti-CD11b) and Mo2 (anti-CD14) derived from ATCC cell lines HB CRL 8026 and TIB 228/HB44. The anti-human gelatinase B antibody Ab110 was raised against a synthetic peptide with the amino acid sequence EALMYPMYRFTEGPPPLHK (SEQ ID NO: 34), which is specific for human gelatinase B (Corcoran, M. L. et al. *J. Biol. Chem.*, 267:515 (1992)).

20 Detection of the immunohistochemical staining was achieved by using a goat anti-rabbit glucose oxidase kit (Vector Laboratories, Burlingame Calif.) according to the manufacturer's directions. Briefly, the sections were rehydrated and pretreated with either acetone or 0.1% trypsin. Normal goat serum was used to block nonspecific binding. Incubation with the primary antibody for 2 hours or overnight (Ab110:1/500 dilution) was followed by either a glucose oxidase labeled secondary anti-rabbit serum, or, in the case of the mouse monoclonal antibodies, were reacted with purified rabbit anti-mouse Ig before incubation with the secondary antibody.

25 Paraffin embedded and frozen sections from osteoclastomas (GCT) were reacted with a rabbit antiserum against gelatinase B (antibody 110) (Corcoran, M. L. et al. *J. Biol. Chem.* 267:515 (1992)), followed by color development with glucose oxidase linked reagents. The osteoclasts of a giant cell tumor were uniformly strongly positive for gelatinase B, whereas the stromal cells were unreactive. Control sections reacted with rabbit preimmune serum were negative. Identical findings were obtained for all 8 long bone giant cell tumors tested (Table IV). The osteoclasts present in three out of four central giant cell granulomas (GCG) of the mandible were also positive for gelatinase B expression. These neoplasms are similar but not identical to the long bone giant cell tumors, apart from their location in the jaws (Shafer, W. G. et al., *Textbook of Oral Pathology*, W. B. Saunders Company, Philadelphia, pp. 144-149 (1983)). In contrast, the multinucleated cells from a peripheral giant cell tumor, which is a generally non-resorptive tumor of oral soft tissue,

14

were unreactive with antibody (Shafer, W. G. et al., *Textbook of Oral Pathology*, W. B. Saunders Company, Philadelphia, pp. 144-149 (1983)).

5 Antibody 110 was also utilized to assess the presence of gelatinase B in normal bone (n=3) and in Paget's disease, in which there is elevated bone remodeling and increased osteoclastic activity. Strong staining for gelatinase B was observed in osteoclasts both in normal bone (mandible of a 2 year old), and in Paget's disease. Staining was again absent in controls incubated with preimmune serum. Osteoblasts did not stain in any of the tissue sections, indicating that gelatinase B expression is limited to osteoclasts in bone. Finally, peripheral blood monocytes were also reactive with antibody 110 (Table IV).

15 TABLE IV

DISTRIBUTION OF GELATINASE B IN VARIOUS TISSUES

Samples	Antibodies tested
	Ab 110
	gelatinase B
GCT frozen (n = 2)	
giant cells	+
stromal cells	-
GCT paraffin (n = 6)	
giant cells	+
stromal cells	-
central GCG (n = 4)	
giant cells	+ (34)
stromal cells	-
peripheral GCT (n = 4)	
giant cells	-
stromal cells	-
Paget's disease (n = 1)	
osteoclasts	+
osteoblasts	-
normal bone (n = 3)	
osteoclasts	+
osteoblasts	-
monocytes (cytopsin)	+

20 Distribution of gelatinase B in multinucleated giant cells, osteoclasts, osteoblasts and stromal cells in various tissues. In general, paraffin embedded tissues were used for these experiments; exceptions are indicated.

Equivalents

25 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 34

5,552,281

15

16

-continued

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 170 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCAAATATCT	AAGTTTATTG	CTTGGATTC	TAGTGAGAGC	TGTTGAATT	GGTGATGTCA	60
AATGTTCTA	GOGTTTTTT	AGTTTGTTT	TATTGAAAAA	TTTAATTATT	TATGCTATAG	120
GTGATATTCT	CTTGAAATAA	ACCTATAATA	GAAAATAGCA	GCAGACAAACA		170

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GTGTCAACCT	GCATATCCTA	AAAATGTCAA	AATGCTGCAT	CTGGTTAACATG	TCGGGGTAGG	60
GGG						63

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTTCCCTCTC	TTGCTTCCCT	TTCCCCAAGCA	GAGGTGCTCA	CTCCATGGCC	ACCGCCACCA	60
CAGGCCACACA	GGGAGTACTG	CCAGACTACT	GCTGATGTTC	TCTTAAGGCC	CAGGGAGTCT	120
CAACCAGCTG	GTGGTGAATG	CTGCCTGGCA	CGGGACCCCC	CCC		163

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 173 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TTTTATTGT	AAATATATGT	ATTACATCCC	TAGAAAAAGA	ATCCCAGGAT	TTTCCCTCCT	60
GTGTGTTTC	GTCTTOCTTC	TTCATGGTCC	ATGATGCCAG	CTGAGGTTGT	CAGTACAATG	120
AAACCAAAC	GGCGGGATG	AAGCAGATTA	TTCTGCCATT	TTTCCAGGTC	TTT	173

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 157 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double

5,552,281

17

18

-continued

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGCTGGACAT GGGTGCCTC CACGTCCTC ATATCCCCAG GCACACTCTG GCCTCAGGTT	60
TTGCCCTGGC CATGTCACT ACCTGGAGTG GGCCCTCCCC TTCTTCAGCC TTGAATCAA	120
AGCCACTTGT TTAGGCAGG ATTCCCAGA CCACTCATCA CATTAAAAAA TATTTTAAA	180
ACAAAAAAA AAAAAAA	197

(2) INFORMATION FOR SEQ ID NO:6

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTGACAAAGC TGTTTATTC CACCAATAAA TAGTATATGG TGATTGGGT TTCTATTTAT	60
AAGAGTAGTG GCTATTATAT GGGGTATCAT GTTGATGCTC ATAAATAGTT CATATCTACT	120
TAATTTGCCT TC	132

(2) INFORMATION FOR SEQ ID NO:7

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 75 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAAGAGAGTT GTATGTACAA CCCAACAGG CAAGGCAGCT AAATGCAGAG GGTACAGAGA	60
GATCCCGAGG GAATT	75

(2) INFORMATION FOR SEQ ID NO:8

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 151 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGATGGAAAC ATGTAGAAAGT CCAGAGAAAA ACAATTTAA AAAAAGGTGG AAAAGTTACG	60
GCAAACCTGA GATTCAGCA TAAAATCTT AGTTAGAAAGT GAGAGAAAGA AGAGGGAGGC	120
TGGTTGCTGT TGCACGTATC AATAGGTTAT C	151

(2) INFORMATION FOR SEQ ID NO:9

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 141 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

5,552,281

19

20

-continued

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TTCTTGATCT TTAGAACACT ATGAATAAGG AAAAAAAGAAA AAACGTGTTCA AAATAAAAATG	60
TAGGAGCCGT GCTTTGGAA TGCTTGAGTG AGGAGCTCAA CAAGTCCTCT CCCAAGAAG	120
CAATGATAAA ACTTGACAAA A	141

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACCCATTCTTCT AACAAATTTT ACTGTAAAAT TTTGGTCAA AGTTCTAACGC TTAATCACAT	60
CTCAAAAGAAT AGAGGCAATA TATAAGCCCAT CTTACTAGAC ATACAGTATT AAACCTGGACT	120
GAATATGAGG ACAAGCTCTA GTGGTCATTA AACCCCTCAG AA	162

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 157 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ACATATATTA ACAGCATTCA TTTGGCCAAA ATCTACACGT TTGTAGAATC CTACTGTATA	60
TAAAGTGGGA ATGTATCAAG TATAGACTAT GAAAGTGCAA ATAACAAGTC AAGGTTAGAT	120
TAACCTTTT TTTTACATT ATAAAATTAA CTTGTTT	157

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 75 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CCAAATTTCT CTGGAATCCA TCCTCCCTCC CATCACCAATA GCCTCGAGAC GTCATTTCTG	60
TTTGACTACT CCAGC	75

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AACTAACCTC CTCOGACCCC TGCCCTCACTC ATTTACACCA ACCACCCAAAC TATCTATAAA	60
CCTGAGCCAT GGCCATCCCT TATGAGCGGC GCAGTGATTA TAGGCTTTCG CTCTAAGATA	120

5,552,281

21

22

-continued

AAAAT

124

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 131 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATTATTATTTC TTTTTTTATG TTAGCTTAGC CATGAAAAAT TTACTGGTGA AGCAGTTAAT	60
AAAAACACACA TCCCATTGAA GGGTTTTGTA CATTTCAGTC CTTACAAATA ACAAAAGCAAT	120
GATAAACCCG GCACGGTCCTG ATAGGAAATT C	151

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 105 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGTGACACAA ACATGCATTC GTTTTATTCA TAAAACAGCC TGGTTTCCTA AAACAATACA	60
AACAGCATGT TCATCAOCAG GAAGCTGGCC GTGGGCAGGG OOOCC	105

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 246 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ATAOGTTAGA TTCTCATTCA CGGGACTAGT TAGCTTTAAG CACCCCTAGAG GACTAGGGTA	60
ATCTGACTTC TCACCTCCTA AGTTCCCTCT TATATCCTCA AGGTAGAAAT GTCTATGTTT	120
TCTACTCCAA TTCATAAAATC TATTCTATAAG TCTTTGGTAC AAOTTACATG ATAAAAAGAA	180
ATGTGATTTC TCTTCCCTTC TTTGCACTTT TGAAATAAAG TATTATCTC CTOTCTACAG	240
TTTAAT	246

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 188 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTCCAGTATA AAGGAAAGCG TTAAGTCGGT AAGCTAGAGG ATTGTAAATA TCTTTTATGT	60
CCTCTAGATA AAACACCCGA TTAACAGATG TTAACCTTT ATGTTTGAT TTGCTTTAAA	120
AATGGCCTTC TACACATTAG CTCCAGCTAA AAAGACACAT TGAGAGCTTA GAGGATAGTC	180

5,552,281

23

24

-continued

TCTGGAGC

188

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 212 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCACTTGAA	GGGAGTTGGT	TGCTTATTT	TGAAOCAGAT	GTGGTGATAAC	TGAGATTGTC	60
TGTTCA	GTTT	CCCCATTGT	TTGTGCTTC	AATGATCCTT	CCTACTTTGC	120
CCATGAC	CTT	TTTCACTGTG	GCCATCAAGG	ACTTTCTGA	CAGCTTGTGT	180
TAAGAGATGT	GA	CTACAGCC	TGCCCC	TGAC	TG	212

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 203 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TGTTAG	TTT	TAGGAAGGCC	TGTCTTCTGG	GAGTGAGGTT	TATTAGTCCA	CTTCTTGAG	60
CTAGAC	GTCC	TATAGTTAGT	CACTGGGAT	GGTGAAGAG	GGAGAAGAGG	AAGGGCGAAG	120
GGAAGGG	GTC	TTTGCTAGTA	TCTCCATTTC	TAGAAGATGG	TTTAGATGAT	AACCACAGGT	180
CTATATGAGC	ATAGTAAGGC	TGT					203

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 177 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CCTATTCTG	ATCCTGACTT	TGGACAAGGC	CCTTCAGCCA	GAAGACTGAC	AAAGTCATCC	60
TCCGTCTACC	AGAGCGTGCA	CTTGTGATCC	AAAAATAAGC	TTCATCTCCG	GCTGTGCCTT	120
GGGTGGAAGG	GGCAGGATTC	TGCAGCTGCT	TTTGCATTTC	TCTTCCTAAA	TTTCATT	177

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 106 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CGGAGCGTAG	GTGTGTTTAT	TCCTGTACAA	ATCATTACAA	AACCAAGTCT	GGGGCAGTCA	60
CCGCCCCCAC	CCATCACCCC	AATGCAATGG	CTAGCTGCTG	GCCTTT		106

5,552,281

25

26

-continued

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 139 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(* i) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TTAGTTCACT CAAACGAGGC AACCCCTTT GGCACGTGCTG CCACGGGGT CATGGCGGTT	60
GTGGCAGCTG CGGAGGTTTC CCCAACACCC TCCCTCTGCTT CCCTGTGTGT CGGGGTCCTCA	120
GGAGCTGACC CAGAGTGGA	139

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 177 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(* i) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GCTGAATGTT TAAGAGAGAT TTTGGTCTTA AAGGCTTCAT CATGAAAGTG TACATGCATA	60
TGCAAGTGTG ATTACGTGG TATGGATGGT TGCTTGTGTTA TTAACTAAAG ATGTACAGCA	120
AACTGCCCGT TTAGAGTCCT CTTAATATTG ATGTCTAAC ACTGGGTCTG CTTATGC	177

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 167 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(* i) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GGCACTGGGA TATGAAATCC AGAAGGGAAA CAAGCACTGG ATAATTAAAA ACAGCTGGGG	60
AGAAAAACTGG OGAAACAAAG GATATATCCT CATGGCTCGA AATAAGAAC ACGCCTGTGG	120
CATTGCCAAC CTGGCCAGCT TCCCCAAGAT GTGACTCCAG CCAGAAA	167

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 151 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(* i) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GCCAGGGCGG ACCGTCCTTA TTCCCTCTCCT GCCTCAGAGG TCAGGAAGGA GGTCTGGCAG	60
GACCTGCACT GGGCCCTAGT CATCTGTGGC AGCGAAGGTG AAGGGACTCA CCTTGTCGCC	120
COTGCCCTGAG TAGAACTTGT TCTGGAATTC C	151

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

5,552,281

27

28

-continued

(A) LENGTH: 156 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AACTCTTCA CACTCTGTA TTTTAGTTT AACAAATATAT GTGTTGTGTC TTGGAAATTA	60
GTCATATCA ATTCAATTG AGCTGTCTCA TTCTTTTTT AATGGTCATA TACAGTGTGA	120
TTCAATTATA AGAATATATC CTAATACTTT TAAAAAA	156

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 150 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GGATAAGAAA GAAGGGCTGA GGCGTAGGGG CGGGGGCTGG CCTGCCTCTC AGTCCTGGGA	60
CGCACAGGCC CGCACAGGTT GAGAGGGCA CTTCCTCTTG CTTAGGTTGG TGAGGATCTG	120
GTCCTGGTTG GCCGGTGGAG AGCCACAAAA	150

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 212 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GCACTTGAA GGGAGTTGGT GTGCTATTTT TGAAGCAGAT GTGGTGATAC TGAGATTGTC	60
TGTTCAAGTTT CCCCATTTGT TTGTGCTTCA AATGATCCTT CCTACTTTGC TTCTCTCCAC	120
CCATGACCTT TTCACTGTG GCCATCAAGG ACTTTCTGA CAGCTTGTGT ACTCTTAGGC	180
TAAGAGATGT GACTACAGCC TGCCCCCTGAC TG	212

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 157 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ATCCCTGGCT GTGGATAGTG CTTTGTGTA GCAAATGCTC CCTCCTTAAG GTTATAAGGC	60
TCCCTGAGTT TGGGAGTGTG GAAGTACTAC TTAACTGTCT GTCCCTGCTTG GCTGTCGTTA	120
TCGTTTCTG GTGATGTTGT OCTAACATA AGAATAC	157

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 152 base pairs
(B) TYPE: nucleic acid

5,552,281

29

30

-continued

(C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GGCTGGGCAT	CCCTCTCCTC	CTCCCATCCCC	ATACATCACC	AGGTCTAATG	TTTACAAACG	60
GTOCCAGCCC	GGCTCTGAAG	CCAAGGGCGG	TCCGTGCCAC	GGTGGCTGTC	AGTATTCCCTC	120
CGTTAGCTT	CCCATAAGGT	TGGAGTATCT	GC			152

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 90 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:31:

CCAACTCCTA	CCCGGATACA	GACCCACAGA	GTGCCATCCC	TGAGAGACCA	GACCGCTCCC	60
CAATACTCTC	CTAAAATAAA	CATGAAGCAC				90

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CATGGATGAA	TGTCTCATGG	TGGGAAGGAA	CATGGTACAT	TTC		43
------------	------------	------------	------------	-----	--	----

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2333 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:33:

AGACACCTCT	GCCCTCACCA	TGAGCCCTTG	GCAGCCCCCTG	GTCCTGGTGC	TCCCTGGTGC	60
GGGCTGCTGC	TTTGCTGCC	CCAGACAGCG	CCAGTCCACC	CTTGTGCTCT	TCCCTGGAGA	120
CCTGAGAAC	AATCTCACCG	ACAGGCAGCT	GGCAGAGGAA	TACCTGTACC	GCTATGGTTA	180
CACTCGGGTG	GCAGAGATGC	GTOGAGAGTC	AAAATCTCTG	GGGCCTGCAC	TGCTGCTTCT	240
CCAGAAGCAA	CTGTCCCTGC	CCOAGACCGG	TOAGCTGGAT	AGCGCCACGC	TGAAGGCCAT	300
GCGAACCCCA	CGGTGCGGGG	TCCCAGACCT	GGGCAGATTC	CAAACCTTTG	A0GGCGACCT	360
CAAGTGGCAC	CACCAACAACA	TCACCTATTG	GATCCAAAAC	TACTCGGAAG	ACTTGCCCGG	420
GGCGGTGATT	GACGACGCC	TTGCCCCGCG	CTTCGCACTG	TGGAGCGCGG	TGACGCCGCT	480
CACCTTCACT	CGCGTGTACA	GCCGGGACCG	AGACATCGTC	ATCCAGTTG	GTOTCGCGGA	540
GCACGGGAGAC	GGGTATCCCT	TCGACGGGAA	GGACGGGCTC	CTGGCACACG	CCTTCCCTCC	600
TGGCCCCGGC	ATTCAGGGAG	ACGCCCCATT	CGACGATGAC	GAGTTGTGGT	CCCTGGGCAA	660

5,552,281

31

32

-continued

GGGCOTCGTG GTTCCAACTC GGTTGGAAA CGCAGATGGC GCGGCCCTGCC ACTTCCCTT	720
CATCTTCGAG GGCGCTCCT ACTCTGCCTG CACCACCGAC GGTCGCTCCG ACGGGTTGCC	780
CTGGTGCAGT ACCACGGCCA ACTACGACAC CGACGACCGG TTGGCTTCT GCCCAAGCGA	840
GAGACTCTAC ACCCGGGACG GCAATGCTGA TGGAAACCC TGCCAGTTTC CATTCATCTT	900
CCAAGGCCAA TCCTACTCCG CCTGCACAC GGACGGTCGC TCCGACGGCT ACCOCTGGTG	960
CGCCACCCACC GCCAACTACG ACCGGGACAA GCTCTCGGC TTCTGCCGA CCCGAGCTGA	1020
CTCGACGCGT ATGGGGGGCGA ACTCGGGGGG GGAGCTGTGC GTCTTCCCCCT TCACCTTCCT	1080
GGGTAAGGAG TACTCGACCT GTACCAGCGA GGGCCGCGGA GATGGGGGCC TCTGOTGCC	1140
TACCACTCG AACTTGACA GCGACAAGAA GTGGGCTTC TGCCCGGCC AAGGATACAG	1200
TTTGTTCCTC GTGGCGGCCG ATGAGTTCGG CCACGGCTG GGCTTAGATC ATTCCCTCAGT	1260
CGCGGAGGCG CTCATGTACC CTATGTACCG TTCACTGAG GGGCCCCCT TGCTATAAGGA	1320
CGACGTGAAT GGCACTCCGC ACCTCTATGG TCCCTGCCCT GAACCTGAGC CACGGCCCTC	1380
AACCACCAAC ACACCGCAGC CCACGGCTCC CCCGACGGTC TGCCCCACCG GACCCCCCAC	1440
TGTCCACCCC TCAGAGEGCC CCACAGCTGG CCCCACAGGT CCCCTCTAG CTGGCCCCAC	1500
AGGTCCCCCCC ACTGCTGGCC CTTCTACGGC CACTACTGTG CCTTGAGTC CGGTGGACGA	1560
TGCCTGCAAC GTGAACATCT TCGACGCCAT CGCGGAGATT GGGAACCGAGC TGTATTTGTT	1620
CAAGGATGGG AAGTACTGGC GATTCTCTOA GGGCAGGGGG AGCCGGCCGC AGGGCCCCCTT	1680
CCTTATCGCC GACAAGTGGC CGCGCTGCC CCGCAAGCTG GACTCGGTCT TTGAGGAGCC	1740
GCTCTCCAAG AAGCTTTCT TCTTCTCTGG CGGCCAGGTG TGGGTGTACA CAGGCGCGTC	1800
GGTGCTGGC CGAGGGCGTC TGGACAAGCT GGGCCTGGGA GCCGACGTGG CCCAGGTGAC	1860
CGGGGCCCTC CGGAGTGGCA GGGGAAGAT GCTGCTGTT AGCGGGCGGC GCCTCTGGAG	1920
GTTCGACGTG AAGGGCGAGA TGGTGGATCC CGGGAGCGCC AGCGAGGTGG ACCGGATGTT	1980
CCCCGGGGTG CCTTGGACA CGCACGACGT CTTCCAGTAC CGAGAGAAAG CCTATTTCTG	2040
CCAGGACCGC TTCTACTGGC GCGTGAGTTC CGGGAGTGAG TTGAACCAGG TOGACCAAGT	2100
GGGCTACGTG ACCTATGACA TCCCTGCAGTG CCCTGAGGAC TAGGGCTCCC GTCCCTGCTTT	2160
GCAGTGCCAT GTAAATCCCC ACTGGGACCA ACCCTGGGG AAGGAGCCAGT TTGCCGGATA	2220
CAAACGGTA TTCTGTTCTG GAGGAAAGGG AGGAGTGGAG GTGGGCTGGG CCCTCTCTTC	2280
TCACCTTTGT TTTTGTGAGTGTCTA ATAAACTTGG ATTCTCTAAC CTTT	2334

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Glu	Ala	Lys	Met	Tyr	Pro	Met	Tyr	Arg	Phe	Thr	Glu	Gly	Pro	Pro	Lys
1															15

His Lys

We claim:

1. An isolated osteoclast-specific or -related DNA sequence, or its complementary sequence, the DNA sequence comprising a nucleic acid sequence selected from the group consisting of:

a) DNA sequences set forth in the group consisting of SEQ ID NOS. 12, 14, 16 and 17, or their complementary strands; and

33

- b) DNA sequences which hybridize under standard conditions to the DNA sequences defined in a).
- 2. A DNA construct capable of replicating, in a host cell, osteoclast-specific or -related DNA, said construct comprising:
 - a) a DNA sequence of claim 1; and
 - b) sequences, in addition to said DNA sequence, necessary for transforming or transfecting a host cell, and for replicating, in a host cell, said DNA sequence.
- 3. A DNA construct capable of replicating and expressing, in a host cell, osteoclast-specific or -related DNA, said construct comprising:

34

- a) a DNA sequence of claim 2; and
- b) sequences, in addition to said DNA sequence, necessary for transforming or transfecting a host cell, and for replicating and expressing, in a host cell, said DNA sequence.
- 4. A cell stably transformed or transfected with a DNA construct according to claim 3.
- 5. A cell stably transformed or transfected with a DNA construct according to claim 4.

* * * * *